

Datasheet: MCA506SBV570

BATCH NUMBER 100008259

Description:	RAT ANTI HUMAN CD235a:StarBright Violet 570
Specificity:	CD235a
Other names:	GLYCOPHORIN A
Format:	StarBright Violet 570
Product Type:	Monoclonal Antibody
Clone:	YTH89.1
Isotype:	IgG2b
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Product Form	Purified IgG conjugat	ed to StarBright Violet	570 - liquid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm
	StarBright Violet 570	404	571
reparation	5		
sparation	Purified IgG prepared supernatant	I by affinity chromatog	raphy on Protein G
•	9		raphy on Protein G
uffer Solution	supernatant	aline	raphy on Protein G
uffer Solution	supernatant Phosphate buffered s	caline (NaN ₃)	raphy on Protein G
uffer Solution reservative tabilisers	Supernatant Phosphate buffered s 0.09% Sodium Azide	caline (NaN ₃)	raphy on Protein G

External	Database
Links	

UniProt:

P02724 Related reagents

Entrez Gene:

2993 GYPA Related reagents

Synonyms

GPA

Fusion Partners

Spleen cells from immunized DA rats were fused with cells of the Y3/Ag.1.2.3.

Specificity

Rat anti Human CD235a antibody, clone YTH89.1 recognizes glycophorin A, a major sialoglycoprotein of the human erythrocyte membrane.

Flow Cytometry

Use 5μ l of the suggested working dilution to label 10^6 cells in 100μ l. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

- 1. Outram, S. *et al.* (1988) Erythromyeloid lineage fidelity is conserved in erythroleukaemia. Leuk Res. 12 (8): 651-7.
- 2. Jokiranta, T.S. & Meri, S. (1993) Biotinylation of monoclonal antibodies prevents their ability to activate the classical pathway of complement. <u>J Immunol</u>. 151 (4): 2124-31.
- 3. Hoang, T. *et al.* (1996) Opposing effects of the basic helix-loop-helix transcription factor SCL on erythroid and monocytic differentiation. <u>Blood. 87: 102-11.</u>
- 4. Babiker, A.A. *et al.* (2002) Transfer of prostasomal CD59 to CD59-deficient red blood cells results in protection against complement-mediated hemolysis. <u>Am J Reprod Immunol. 47 (3): 183-92.</u>
- 5. Lahlil, R. *et al.* (2004) SCL assembles a multifactorial complex that determines glycophorin A expression. <u>Mol Cell Biol. 24: 1439-52.</u>
- 6. Challier, J.C. *et al.* (2005) Immunocytological evidence for hematopoiesis in the early human placenta. <u>Placenta. 26: 282-8.</u>
- 7. Huang, Y.C. *et al.* (2009) Oral small-molecule tyrosine kinase inhibitor midostaurin (PKC412) inhibits growth and induces megakaryocytic differentiation in human leukemia cells. <u>Toxicol In Vitro. 23: 979-85.</u>
- 8. Tiziani, S. *et al.* (2009) Metabolomic profiling of drug responses in acute myeloid leukaemia cell lines. <u>PLoS One. 2009;4(1):e4251.</u>
- 9. Basu, S. (2010) Erythrocyte membrane defects and asymmetry in paroxysmal nocturnal hemoglobinuria and myelodysplastic syndrome. <u>Hematology. 15: 236-9.</u>
- 10. Saha, S. *et al.* (2011) Elevated levels of redox regulators, membrane-bound globin chains, and cytoskeletal protein fragments in hereditary spherocytosis erythrocyte proteome. <u>Eur J Haematol. 87: 259-66.</u>
- 11. Lucky AB *et al.* (2016) Plasmodium knowlesi Skeleton-Binding Protein 1 Localizes to the 'Sinton and Mulligan' Stipplings in the Cytoplasm of Monkey and Human Erythrocytes. PLoS One. 11 (10): e0164272.
- 12. Scanlon, V.M. *et al.* (2022) Multiparameter analysis of timelapse imaging reveals kinetics of megakaryocytic erythroid progenitor clonal expansion and differentiation. <u>Sci Rep. 12 (1): 16218.</u>

Storage	Store at +4°C. DO NOT FREEZE.
	This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety	Material Safety Datasheet documentation #20471 available at:
Information	https://www.bio-rad-antibodies.com/SDS/MCA506SBV570
	20471
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M426144:231121'

Printed on 17 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint