

Datasheet: MCA519G

BATCH NUMBER 163611

Description:	RAT ANTI MOUSE MACROPHAGES/MONOCYTES
Specificity:	MACROPHAGES/MONOCYTES
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MOMA-2
Isotype:	IgG2b
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			
Immunohistology - Frozen	▪			1/25
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code [BUF09](#)) is recommended for this purpose.

Target Species	Mouse
Product Form	Purified IgG - liquid.
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline.
Preservative	<0.1% Sodium Azide (NaN ₃).

Stabilisers

Approx. Protein Concentrations IgG concentration 0.5 mg/ml.

Immunogen Mouse lymph node stroma.

RRID AB_321970

Fusion Partners Spleen cells from immunized Wistar rats were fused with cells of the SP/0 myeloma cell line.

Specificity **Rat anti Mouse Macrophages/Monocytes antibody, clone MOMA-2** recognizes an intracellular antigen of mouse macrophages and monocytes. It reacts strongly with macrophages in lymphoid organs such as tingible body macrophages and macrophages in T cell dependant areas and is extremely useful in immunohistochemistry. Reacts on all mouse strains tested.

References

1. Gal, A. *et al.* (1996) Nitric oxide production in SJL mice bearing the RcsX lymphoma: a model for in vivo toxicological evaluation of NO. [Proc Natl Acad Sci U S A. 93 \(21\): 11499-503.](#)
2. Dieleman, L.A. *et al.* (1998) Chronic experimental colitis induced by dextran sulphate sodium (DSS) is characterized by Th1 and Th2 cytokines. [Clin Exp Immunol. 114: 385-91.](#)
3. Fischer, M.B. *et al.* (1998) Local synthesis of C3 within the splenic lymphoid compartment can reconstitute the impaired immune response in C3-deficient mice. [J Immunol. 160: 2619-25.](#)
4. Jiang, H.R. *et al.* (1999) Macrophages and dendritic cells in IRBP-induced experimental autoimmune uveoretinitis in B10RIII mice. [Invest Ophthalmol Vis Sci. 40: 3177-85.](#)
5. Babaei, S. *et al.* (2000) Blockade of endothelin receptors markedly reduces atherosclerosis in LDL receptor deficient mice: role of endothelin in macrophage foam cell formation. [Cardiovasc Res. 2000 Oct;48: 158-67.](#)
6. Oguchi, S. *et al.* (2000) Monoclonal antibody against vascular cell adhesion molecule-1 inhibits neointimal formation after periadventitial carotid artery injury in genetically hypercholesterolemic mice. [Arterioscler Thromb Vasc Biol. 20: 1729-36.](#)
7. Kusunoki, J. *et al.* (2001) Acyl-CoA:cholesterol acyltransferase inhibition reduces atherosclerosis in apolipoprotein E-deficient mice. [Circulation. 103: 2604-9.](#)
8. Lesnik, P. *et al.* (2003) Decreased atherosclerosis in CX3CR1-/- mice reveals a role for fractalkine in atherogenesis. [J Clin Invest. 111: 333-40.](#)
9. Oguro, A. *et al.* (2003) NaF induces early differentiation of murine bone marrow cells along the granulocytic pathway but not the monocytic or preosteoclastic pathway *in vitro*. [In Vitro Cell Dev Biol Anim. 39 \(5-6\): 243-8.](#)
10. Yun, J.J. *et al.* (2004) Combined blockade of the chemokine receptors CCR1 and CCR5 attenuates chronic rejection. [Circulation. 109: 932-7.](#)
11. Addison, C.L. *et al.* (2004) Overexpression of the duffy antigen receptor for chemokines (DARC) by NSCLC tumor cells results in increased tumor necrosis. [BMC Cancer. 4: 28.](#)
12. Nakai, Y. *et al.* (2004) Natural killer T cells accelerate atherogenesis in mice. [Blood. 104 \(7\): 2051-9.](#)

13. Persson, L. *et al.* (2005) Immunoglobulin treatment reduces atherosclerosis in apolipoprotein E^{-/-} low-density lipoprotein receptor^{-/-} mice via the complement system. [Clin Exp Immunol. 142: 441-5.](#)
14. Wang, H. *et al.* (2006) Activated macrophages are essential in a murine model for T cell-mediated chronic psoriasiform skin inflammation. [J Clin Invest. 116: 2105-14.](#)
15. Bourdillon, M.C. *et al.* (2006) Reduced atherosclerotic lesion size in P-selectin deficient apolipoprotein E-knockout mice fed a chow but not a fat diet. [J Biomed Biotechnol. 2006 \(2\): 49193.](#)
16. Bhatia, V.K. *et al.* (2007) Complement C1q reduces early atherosclerosis in low-density lipoprotein receptor-deficient mice. [Am J Pathol. 170: 416-26.](#)
17. Nix, R.N. *et al.* (2007) Hemophagocytic macrophages harbor *Salmonella enterica* during persistent infection. [PLoS Pathog. 3\(12\):e193.](#)
18. Wang, Z. *et al.* (2008) Phospholipase C beta3 deficiency leads to macrophage hypersensitivity to apoptotic induction and reduction of atherosclerosis in mice. [J Clin Invest. 118: 195-204.](#)
19. Westmuckett, A.D. *et al.* (2009) Lack of tyrosylprotein sulfotransferase activity in hematopoietic cells drastically attenuates atherosclerosis in Ldlr^{-/-} mice. [Arterioscler Thromb Vasc Biol. 29: 1730-6.](#)
20. Leung, V.W. *et al.* (2009) Decay-accelerating factor suppresses complement C3 activation and retards atherosclerosis in low-density lipoprotein receptor-deficient mice. [Am J Pathol. 175: 1757-67.](#)
21. Yuan, Z. *et al.* (2009) Quantitative trait locus analysis of neointimal formation in an intercross between C57BL/6 and C3H/HeJ apolipoprotein E-deficient mice. [Circ Cardiovasc Genet. 2: 220-8.](#)
22. Pillai, M.M. *et al.* (2009) Inducible transgenes under the control of the hCD68 promoter identifies mouse macrophages with a distribution that differs from the F4/80 - and CSF-1R-expressing populations. [Exp Hematol. 37: 1387-92.](#)
23. Pedersen, T.X. *et al.* (2010) The pro-inflammatory effect of uraemia overrules the anti-atherogenic potential of immunization with oxidized LDL in apoE^{-/-} mice. [Nephrol Dial Transplant. 25: 2486-91.](#)
24. Chen, S. (2010) IL-17A is proatherogenic in high-fat diet-induced and *Chlamydia pneumoniae* infection-accelerated atherosclerosis in mice. [J Immunol. 185: 5619-27.](#)
25. Gao, Q. *et al.* (2010) A critical function of Th17 proinflammatory cells in the development of atherosclerotic plaque in mice. [J Immunol. 185: 5820-7.](#)
26. Madrigal-Matute, J. *et al.* (2010) Heat shock protein 90 inhibitors attenuate inflammatory responses in atherosclerosis. [Cardiovasc Res. 86 \(2\): 330-7.](#)
27. Duewell, P. *et al.* (2010) NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. [Nature. 464: 1357-61.](#)
28. Li, J. *et al.* (2011) Cardiac tissue-restricted deletion of plakoglobin results in progressive cardiomyopathy and activation of {beta}-catenin signaling. [Mol Cell Biol. 31: 1134-44.](#)
29. Skoura, A. *et al.* (2011) Sphingosine-1-phosphate receptor-2 function in myeloid cells regulates vascular inflammation and atherosclerosis. [Arterioscler Thromb Vasc Biol. 31 \(1\): 81-5.](#)
30. Frossard, J.L. *et al.* (2011) Role of CCL-2, CCR-2 and CCR-4 in cerulein-induced acute pancreatitis and pancreatitis-associated lung injury. [J Clin Pathol. 64 \(5\): 387-93.](#)
31. de Jager, S.C. *et al.* (2011) Growth differentiation factor 15 deficiency protects against

- atherosclerosis by attenuating CCR2-mediated macrophage chemotaxis. [J Exp Med. 208 \(2\): 217-25.](#)
32. Ng, H.P. *et al.* (2011) Attenuated atherosclerotic lesions in apoE-Fcy-chain-deficient hyperlipidemic mouse model is associated with inhibition of Th17 cells and promotion of regulatory T cells. [J Immunol. 187 \(11\): 6082-93.](#)
33. Weingärtner, O. *et al.* (2011) Differential effects on inhibition of cholesterol absorption by plant stanol and plant sterol esters in apoE-/- mice. [Cardiovasc Res. 90: 484-92.](#)
34. Yamamoto, S. *et al.* (2011) Oral activated charcoal adsorbent (AST-120) ameliorates extent and instability of atherosclerosis accelerated by kidney disease in apolipoprotein E-deficient mice. [Nephrol Dial Transplant. 26 \(8\): 2491-7.](#)
35. Che, J. *et al.* (2011) Endothelial FGF receptor signaling accelerates atherosclerosis. [Am J Physiol Heart Circ Physiol. 300: H154-61.](#)
36. Ruf, M.T. *et al.* (2012) Chemotherapy-Associated Changes of Histopathological Features of *Mycobacterium ulcerans* Lesions in a Buruli Ulcer Mouse Model. [Antimicrob Agents Chemother. 56: 687-96.](#)
37. van der Sluis, R.J. *et al.* (2014) Prolactin receptor antagonism uncouples lipids from atherosclerosis susceptibility. [J Endocrinol. 222 \(3\): 341-50.](#)
38. Lee, M.R. *et al.* (2014) The adipokine Retnla modulates cholesterol homeostasis in hyperlipidemic mice. [Nat Commun. 5: 4410.](#)
39. Song, G. *et al.* (2015) Molecular hydrogen stabilizes atherosclerotic plaque in low-density lipoprotein receptor-knockout mice. [Free Radic Biol Med. 87: 58-68.](#)
40. Shuto, Y. *et al.* (2015) Repetitive Glucose Spikes Accelerate Atherosclerotic Lesion Formation in C57BL/6 Mice. [PLoS One. 10 \(8\): e0136840.](#)
41. Takata, H. *et al.* (2015) Vascular angiotensin II type 2 receptor attenuates atherosclerosis via a kinin/NO-dependent mechanism. [J Renin Angiotensin Aldosterone Syst. 16 \(2\): 311-20.](#)
42. van der Sluis, R.J. *et al.* (2015) Haloperidol inhibits the development of atherosclerotic lesions in LDL receptor knockout mice. [Br J Pharmacol. 172 \(9\): 2397-405.](#)
43. Wan W *et al.* (2015) Atypical chemokine receptor 1 deficiency reduces atherogenesis in ApoE-knockout mice. [Cardiovasc Res. 106 \(3\): 478-87.](#)
44. Oguiza A *et al.* (2015) Peptide-based inhibition of IκB kinase/nuclear factor-κB pathway protects against diabetes-associated nephropathy and atherosclerosis in a mouse model of type 1 diabetes. [Diabetologia. 58 \(7\): 1656-67.](#)
45. Yamamoto, S. *et al.* (2015) Atherosclerosis following renal injury is ameliorated by pioglitazone and losartan via macrophage phenotype [Atherosclerosis. 242 \(1\): 56-64.](#)
46. Krishack, P.A. *et al.* (2015) Serum Amyloid A Facilitates Early Lesion Development in Ldlr-/- Mice. [J Am Heart Assoc. 4 \(7\): pii: e001858.](#)
47. Aoki, S. *et al.* (2015) Oral administration of the β-glucan produced by *Aureobasidium pullulans* ameliorates development of atherosclerosis in apolipoprotein E deficient mice [Journal of Functional Foods. 18: 22-7.](#)
48. Peng, Y. *et al.* (2016) Inactivation of Semicarbazide-Sensitive Amine Oxidase Stabilizes the Established Atherosclerotic Lesions via Inducing the Phenotypic Switch of Smooth Muscle Cells. [PLoS One. 11 \(4\): e0152758.](#)
49. Grootaert, M.O. *et al.* (2016) NecroX-7 reduces necrotic core formation in atherosclerotic plaques of Apoe knockout mice. [Atherosclerosis. 252: 166-74.](#)
50. Hong, Y.F. *et al.* (2016) *Lactobacillus acidophilus* K301 Inhibits Atherogenesis via Induction of 24 (S), 25-Epoxycholesterol-Mediated ABCA1 and ABCG1 Production and

- Cholesterol Efflux in Macrophages. [PLoS One. 11 \(4\): e0154302.](#)
51. Bosteen, M.H. *et al.* (2017) Effects of apolipoprotein M in uremic atherosclerosis. [Atherosclerosis. 265: 93-101.](#)
52. Neele, A.E. *et al.* (2018) Myeloid Kdm6b deficiency results in advanced atherosclerosis. [Atherosclerosis. 275: 156-65.](#)
53. Luque-Martin, R. *et al.* (2019) Targeting Histone Deacetylases in Myeloid Cells Inhibits Their Maturation and Inflammatory Function With Limited Effects on Atherosclerosis. [Front Pharmacol. 10: 1242.](#)
54. Sun, Y. *et al.* (2019) Silencing of junctional adhesion molecule-like protein attenuates atherogenesis and enhances plaque stability in ApoE^{-/-} mice. [Clin Sci \(Lond\). 133 \(11\): 1215-28.](#)
55. Tang, G. *et al.* (2019) Metformin inhibited Nod-like receptor protein 3 inflammasomes activation and suppressed diabetes-accelerated atherosclerosis in apoE^{-/-} mice. [Biomed Pharmacother. 119: 109410.](#)
56. van Duijn, J. *et al.* (2019) CD8+ T-cells contribute to lesion stabilization in advanced atherosclerosis by limiting macrophage content and CD4+ T-cell responses. [Cardiovasc Res. 115 \(4\): 729-38.](#)
57. Zhao, J. *et al.* (2019) Atherogenesis in the Carotid Artery with and without Interrupted Blood Flow of Two Hyperlipidemic Mouse Strains. [J Vasc Res. 56 \(5\): 241-254.](#)
58. Burris, R.L. *et al.* (2020) Maternal exposure to soy diet reduces atheroma in hyperlipidemic F1 offspring mice by promoting macrophage and T cell anti-inflammatory responses. [Atherosclerosis. 313: 26-34.](#)
59. Baardman, J. *et al.* (2020) Macrophage ATP citrate lyase deficiency stabilizes atherosclerotic plaques. [Nat Commun. 11 \(1\): 6296.](#)
60. Kato, T. *et al.* (2020) A non-selective endothelin receptor antagonist bosentan modulates kinetics of bone marrow-derived cells in ameliorating pulmonary hypertension in mice. [Pulm Circ. 10 \(2\): 2045894020919355.](#)
61. Douna, H. *et al.* (2020) B- and T-lymphocyte attenuator stimulation protects against atherosclerosis by regulating follicular B cells. [Cardiovasc Res. 116 \(2\): 295-305.](#)
62. Gajeton, J. *et al.* (2020) miR-467 Prevents Inflammation And Insulin Resistance In Response To Hyperglycemia [BioRxiv. Jun 11 Preprint \[Epub ahead of print\].](#)
63. Lecoeur, H. *et al.* (2020) Targeting Macrophage Histone H3 Modification as a Leishmania Strategy to Dampen the NF-κB/NLRP3-Mediated Inflammatory Response. [Cell Rep. 30 \(6\): 1870-1882.e4.](#)
64. Gajeton, J. *et al.* (2021) Hyperglycemia-Induced miR-467 Drives Tumor Inflammation and Growth in Breast Cancer. [Cancers \(Basel\). 13 \(6\): 1346.](#)
65. Zhao, J. *et al.* (2021) Inflammation and enhanced atherogenesis in the carotid artery with altered blood flow in an atherosclerosis-resistant mouse strain. [Physiol Rep. 9 \(11\): e14829.](#)
66. Zhao, J. *et al.* (2021) Inflammation and enhanced atherogenesis in the carotid artery with altered blood flow in an atherosclerosis-resistant mouse strain. [Physiol Rep. 9 \(11\): e14829.](#)
67. van Duijn, J. *et al.* (2021) Tc17 CD8+ T cells accumulate in murine atherosclerotic lesions, but do not contribute to early atherosclerosis development. [Cardiovasc Res. 117 \(14\): 2755-66.](#)
68. Imai, T. *et al.* (2022) Smooth muscle cell specific NEMO deficiency inhibits atherosclerosis in ApoE^{-/-} mice. [Sci Rep. 12 \(1\): 12538.](#)

69. Smit, V. *et al.* (2023) Single-cell profiling reveals age-associated immunity in atherosclerosis. [Cardiovasc Res. 119 \(15\): 2508-21.](#)
70. Karasawa, T. *et al.* (2023) Caspase-11 deficiency attenuates neutrophil recruitment into the atherosclerotic lesion in apolipoprotein E-deficient mice. [Biochem Biophys Res Commun. 686: 149158.](#)
71. Segers, F.M.E. *et al.* (2023) Magnetic resonance imaging contrast-enhancement with superparamagnetic iron oxide nanoparticles amplifies macrophage foam cell apoptosis in human and murine atherosclerosis. [Cardiovasc Res. 118 \(17\): 3346-59.](#)
72. Grievink, H.W. *et al.* (2021) Stimulation of the PD-1 Pathway Decreases Atherosclerotic Lesion Development in Ldlr Deficient Mice. [Front Cardiovasc Med. 8: 740531.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA519G>
10040

Regulatory For research purposes only.

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M404314:220822'

Printed on 11 Apr 2024