

Datasheet: MCA519G

Description:	RAT ANTI MOUSE MACROPHAGES/MONOCYTES
Specificity:	MACROPHAGES/MONOCYTES
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MOMA-2
Isotype:	IgG2b
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			
Immunohistology - Frozen	▪			1/25
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

*Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.

(1)Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
Immunogen	Mouse lymph node stroma.
RRID	AB_321970
Fusion Partners	Spleen cells from immunised Wistar rats were fused with cells of the SP/0 myeloma cell line.
Specificity	Rat anti Mouse Macrophages/Monocytes antibody, clone MOMA-2 recognizes an intracellular antigen of mouse macrophages and monocytes. It reacts strongly with macrophages in lymphoid organs such as tingible body macrophages and macrophages in T cell dependant areas and is extremely useful in immunohistochemistry. Reacts on all mouse strains tested.
References	<ol style="list-style-type: none"> 1. van der Sluis, R.J. <i>et al.</i> (2014) Prolactin receptor antagonism uncouples lipids from atherosclerosis susceptibility. J Endocrinol. 222 (3): 341-50. 2. Nakai, Y. <i>et al.</i> (2004) Natural killer T cells accelerate atherogenesis in mice. Blood. 104 (7): 2051-9. 3. Skoura, A. <i>et al.</i> (2011) Sphingosine-1-phosphate receptor-2 function in myeloid cells regulates vascular inflammation and atherosclerosis. Arterioscler Thromb Vasc Biol. 31 (1): 81-5. 4. Madrigal-Matute, J. <i>et al.</i> (2010) Heat shock protein 90 inhibitors attenuate inflammatory responses in atherosclerosis. Cardiovasc Res. 86 (2): 330-7. 5. de Jager, S.C. <i>et al.</i> (2011) Growth differentiation factor 15 deficiency protects against atherosclerosis by attenuating CCR2-mediated macrophage chemotaxis. J Exp Med. 208 (2): 217-25. 6. Frossard, J.L. <i>et al.</i> (2011) Role of CCL-2, CCR-2 and CCR-4 in cerulein-induced acute pancreatitis and pancreatitis-associated lung injury. J Clin Pathol. 64 (5): 387-93. 7. Bhatia, V.K. <i>et al.</i> (2007) Complement C1q reduces early atherosclerosis in low-density lipoprotein receptor-deficient mice. Am J Pathol. 170: 416-26. 8. Bourdillon, M.C. <i>et al.</i> (2006) Reduced atherosclerotic lesion size in P-selectin deficient apolipoprotein E-knockout mice fed a chow but not a fat diet. J Biomed Biotechnol. 2006 (2): 49193. 9. Duewell, P. <i>et al.</i> (2010) NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. Nature. 464: 1357-61. 10. Weingärtner, O. <i>et al.</i> (2011) Differential effects on inhibition of cholesterol absorption by plant stanol and plant sterol esters in apoE^{-/-} mice. Cardiovasc Res. 90: 484-92. 11. Yamamoto, S. <i>et al.</i> (2011) Oral activated charcoal adsorbent (AST-120) ameliorates extent and instability of atherosclerosis accelerated by kidney disease in apolipoprotein E-deficient mice. Nephrol Dial Transplant. 26 (8): 2491-7. 12. Ng, H.P. <i>et al.</i> (2011) Attenuated atherosclerotic lesions in apoE-Fcy-chain-deficient hyperlipidemic mouse model is associated with inhibition of Th17 cells and promotion of regulatory T cells. J Immunol. 187 (11): 6082-93.

13. Ruf, M.T. *et al.* (2012) Chemotherapy-Associated Changes of Histopathological Features of *Mycobacterium ulcerans* Lesions in a Buruli Ulcer Mouse Model. [Antimicrob Agents Chemother. 56: 687-96.](#)
14. Che, J. *et al.* (2011) Endothelial FGF receptor signaling accelerates atherosclerosis. [Am J Physiol Heart Circ Physiol. 300: H154-61.](#)
15. Chen, S. (2010) IL-17A is proatherogenic in high-fat diet-induced and *Chlamydia pneumoniae* infection-accelerated atherosclerosis in mice. [J Immunol. 185: 5619-27.](#)
16. Dieleman, L.A. *et al.* (1998) Chronic experimental colitis induced by dextran sulphate sodium (DSS) is characterized by Th1 and Th2 cytokines. [Clin Exp Immunol. 114: 385-91.](#)
17. Gao, Q. *et al.* (2010) A critical function of Th17 proinflammatory cells in the development of atherosclerotic plaque in mice. [J Immunol. 185: 5820-7.](#)
18. Pedersen, T.X. *et al.* (2010) The pro-inflammatory effect of uraemia overrules the anti-atherogenic potential of immunization with oxidized LDL in apoE^{-/-} mice. [Nephrol Dial Transplant. 25: 2486-91.](#)
19. Lee, M.R. *et al.* (2014) The adipokine Retnla modulates cholesterol homeostasis in hyperlipidemic mice. [Nat Commun. 5: 4410.](#)
20. Hoeksema, M.A. *et al.* (2014) Targeting macrophage Histone deacetylase 3 stabilizes atherosclerotic lesions. [EMBO Mol Med. 6 \(9\): 1124-32.](#)
21. Yamamoto, S. *et al.* (2015) Atherosclerosis following renal injury is ameliorated by pioglitazone and losartan via macrophage phenotype [Atherosclerosis. 242 \(1\): 56-64.](#)
22. Wan W *et al.* (2015) Atypical chemokine receptor 1 deficiency reduces atherogenesis in ApoE-knockout mice. [Cardiovasc Res. 106 \(3\): 478-87.](#)
23. Babaei, S. *et al.* (2000) Blockade of endothelin receptors markedly reduces atherosclerosis in LDL receptor deficient mice: role of endothelin in macrophage foam cell formation. [Cardiovasc Res. 2000 Oct;48: 158-67.](#)
24. Krishack, P.A. *et al.* (2015) Serum Amyloid A Facilitates Early Lesion Development in Ldlr^{-/-} Mice. [J Am Heart Assoc. 4 \(7\): pii: e001858.](#)
25. Aoki, S. *et al.* (2015) Oral administration of the β -glucan produced by *Aureobasidium pullulans* ameliorates development of atherosclerosis in apolipoprotein E deficient mice [Journal Funct Foods. 18: 22-7.](#)
26. Song, G. *et al.* (2015) Molecular hydrogen stabilizes atherosclerotic plaque in low-density lipoprotein receptor-knockout mice. [Free Radic Biol Med. 87: 58-68.](#)
27. Takata, H. *et al.* (2015) Vascular angiotensin II type 2 receptor attenuates atherosclerosis via a kinin/NO-dependent mechanism. [J Renin Angiotensin Aldosterone Syst. 16 \(2\): 311-20.](#)
28. Wezel, A. *et al.* (2015) Mast cells mediate neutrophil recruitment during atherosclerotic plaque progression. [Atherosclerosis. 241 \(2\): 289-96.](#)
29. Oguiza A *et al.* (2015) Peptide-based inhibition of I κ B kinase/nuclear factor- κ B pathway protects against diabetes-associated nephropathy and atherosclerosis in a mouse model of type 1 diabetes. [Diabetologia. 58 \(7\): 1656-67.](#)
30. Shuto, Y. *et al.* (2015) Repetitive Glucose Spikes Accelerate Atherosclerotic Lesion Formation in C57BL/6 Mice. [PLoS One. 10 \(8\): e0136840.](#)
31. Peng, Y. *et al.* (2016) Inactivation of Semicarbazide-Sensitive Amine Oxidase Stabilizes the Established Atherosclerotic Lesions via Inducing the Phenotypic Switch of Smooth Muscle Cells. [PLoS One. 11 \(4\): e0152758.](#)
32. Hong, Y.F. *et al.* (2016) *Lactobacillus acidophilus* K301 Inhibits Atherogenesis via Induction of 24 (S), 25-Epoxycholesterol-Mediated ABCA1 and ABCG1 Production and

- Cholesterol Efflux in Macrophages. [PLoS One. 11 \(4\): e0154302.](#)
33. Grootaert, M.O. *et al.* (2016) NecroX-7 reduces necrotic core formation in atherosclerotic plaques of Apoe knockout mice. [Atherosclerosis. 252: 166-74.](#)
34. Oguro, A. *et al.* (2003) NaF induces early differentiation of murine bone marrow cells along the granulocytic pathway but not the monocytic or preosteoclastic pathway *in vitro*. [In Vitro Cell Dev Biol Anim. 39 \(5-6\): 243-8.](#)
35. van der Sluis, R.J. *et al.* (2015) Haloperidol inhibits the development of atherosclerotic lesions in LDL receptor knockout mice. [Br J Pharmacol. 172 \(9\): 2397-405.](#)
36. Addison, C.L. *et al.* (2004) Overexpression of the duffy antigen receptor for chemokines (DARC) by NSCLC tumor cells results in increased tumor necrosis. [BMC Cancer. 4: 28.](#)
37. Neele, A.E. *et al.* (2018) Myeloid Kdm6b deficiency results in advanced atherosclerosis. [Atherosclerosis. 275: 156-65.](#)
38. Tang, G. *et al.* (2019) Metformin inhibited Nod-like receptor protein 3 inflammasomes activation and suppressed diabetes-accelerated atherosclerosis in apoE^{-/-} mice. [Biomed Pharmacother. 119: 109410.](#)
39. van Duijn, J. *et al.* (2019) CD8⁺ T-cells contribute to lesion stabilization in advanced atherosclerosis by limiting macrophage content and CD4⁺ T-cell responses. [Cardiovasc Res. 115 \(4\): 729-38.](#)
40. Kato, T. *et al.* (2020) A non-selective endothelin receptor antagonist bosentan modulates kinetics of bone marrow-derived cells in ameliorating pulmonary hypertension in mice. [Pulm Circ. 10 \(2\): 2045894020919355.](#)
41. Gajeton, J. *et al.* (2020) miR-467 Prevents Inflammation And Insulin Resistance In Response To Hyperglycemia [BioRxiv. Jun 11 Preprint \[Epub ahead of print\].](#)
42. Lecoeur, H. *et al.* (2020) Targeting Macrophage Histone H3 Modification as a Leishmania Strategy to Dampen the NF-κB/NLRP3-Mediated Inflammatory Response. [Cell Rep. 30 \(6\): 1870-1882.e4.](#)
43. van Duijn, J. *et al.* (2020) Tc17 CD8⁺ T cells accumulate in murine atherosclerotic lesions, but do not contribute to early atherosclerosis development [Cardiovasc Res. Oct 16 \[Epub ahead of print\].](#)
44. Baardman, J. *et al.* (2020) Macrophage ATP citrate lyase deficiency stabilizes atherosclerotic plaques. [Nat Commun. 11 \(1\): 6296.](#)
45. Luque-Martin, R. *et al.* (2019) Targeting Histone Deacetylases in Myeloid Cells Inhibits Their Maturation and Inflammatory Function With Limited Effects on Atherosclerosis. [Front Pharmacol. 10: 1242.](#)
46. Zhao, J. *et al.* (2019) Atherogenesis in the Carotid Artery with and without Interrupted Blood Flow of Two Hyperlipidemic Mouse Strains. [J Vasc Res. 56 \(5\): 241-254.](#)
47. Burris, R.L. *et al.* (2020) Maternal exposure to soy diet reduces atheroma in hyperlipidemic F1 offspring mice by promoting macrophage and T cell anti-inflammatory responses. [Atherosclerosis. 313: 26-34.](#)
48. Douna, H. *et al.* (2020) B- and T-lymphocyte attenuator stimulation protects against atherosclerosis by regulating follicular B cells. [Cardiovasc Res. 116 \(2\): 295-305.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR16...)	DyLight@800
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight@650 , DyLight@800
Goat Anti Rat IgG (STAR72...)	HRP
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M389573:210806'

Printed on 29 Aug 2021