

## Datasheet: MCA52FA

<b>Description:</b>	MOUSE ANTI RAT CD5:FITC
<b>Specificity:</b>	CD5
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OX-19
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	50 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Rat		
<b>Product Form</b>	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	FITC	490	525
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A		
<b>Buffer Solution</b>	Phosphate buffered saline		
<b>Preservative</b>	0.09% Sodium Azide		
<b>Stabilisers</b>	1% Bovine Serum Albumin		
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml		
<b>Immunogen</b>	Rat thymocyte glycoproteins.		

External Database  
Links

UniProt:

[P51882](#)    [Related reagents](#)

Entrez Gene:

[54236](#)    Cd5    [Related reagents](#)

---

RRID

AB\_2075315

---

Fusion Partners

Spleen cells from an immunized BALB/c mouse were fused with cells from the NS1 mouse myeloma cell line.

---

Specificity

**Mouse anti Rat CD5 antibody, clone OX-19** recognizes the rat CD5 cell surface antigen, a 69kD glycoprotein expressed by T cells, thymocytes and a subset of B cells.

Mouse anti Rat CD5 antibody, clone OX-19 has been reported as being suitable for use on periodate-lysine paraformaldehyde (PLP) fixed paraffin embedded tissue ([Whiteland et al. 1995](#)).

Mouse anti Rat CD5 antibody, clone OX-19 is routinely tested in flow cytometry on rat splenocytes.

---

Flow Cytometry

Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.

---

References

1. Dallman, M.J. *et al.* (1982) The roles of host and donor cells in the rejection of skin allografts by T cell-deprived rats injected with syngeneic T cells. [Eur J Immunol. 12 \(6\): 511-8.](#)
2. Dallman, M.J. *et al.* (1984) MRC OX-19: a monoclonal antibody that labels rat T lymphocytes and augments in vitro proliferative responses. [Eur J Immunol. 14 \(3\): 260-7.](#)
3. Huitinga, I. *et al.* (1995) Macrophages in T cell line-mediated, demyelinating, and chronic relapsing experimental autoimmune encephalomyelitis in Lewis rats. [Clin Exp Immunol. 100 \(2\): 344-51.](#)
4. Whiteland, J.L. *et al.* (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. [J Histochem Cytochem. 43 \(3\): 313-20.](#)
5. Song, E. *et al.* (2002) Early application of Met-RANTES ameliorates chronic allograft nephropathy. [Kidney Int. 61: 676-85.](#)
6. Moreno-Manzano, V. *et al.* (2003) Retinoids as a potential treatment for experimental puromycin-induced nephrosis [Br J Pharmacol. 139: 823-31.](#)
7. Shiraishi, H. *et al.* (2002) Antibody binding to fas ligand attenuates inflammatory cell infiltration and cytokine secretion, leading to reduction of myocardial infarct areas and reperfusion injury. [Lab Invest. 82: 1121-9.](#)
8. Rusai, K. *et al.* (2008) Administration of interleukin-1 receptor antagonist ameliorates renal ischemia-reperfusion injury. [Transpl Int. 21: 572-80.](#)
9. Antus, B. *et al.* (2001) Contribution of androgens to chronic allograft nephropathy is mediated by dihydrotestosterone [Kidney Int. 60: 1955-63.](#)
10. Antus, B. *et al.* (2005) Effects of progesterone and selective oestrogen receptor modulators on chronic allograft nephropathy in rats. [Nephrol Dial Transplant. 20: 329-35.](#)

11. Kikuchi, H. *et al.* (2000) Severe proteinuria, sustained for 6 months, induces tubular epithelial cell injury and cell infiltration in rats but not progressive interstitial fibrosis. [Nephrol Dial Transplant.15: 799-810.](#)
12. Abrams, M.B. *et al.* (2009) Multipotent mesenchymal stromal cells attenuate chronic inflammation and injury-induced sensitivity to mechanical stimuli in experimental spinal cord injury. [Restor Neurol Neurosci. 27: 307-21.](#)
13. Robichon, R. *et al.* (2005) Pig xenografts to the immunocompetent rat brain: Survival rates using distinct neurotoxic lesions in the nigrostriatal pathway and two rat strains. [Exp Neurol. 194: 333-40.](#)
14. de Vries, H.E. *et al.* (2002) Signal-regulatory protein alpha-CD47 interactions are required for the transmigration of monocytes across cerebral endothelium. [J Immunol. 168: 5832-9.](#)
15. Pamukcu, O. *et al.* (2016) Anti-inflammatory role of obestatin in autoimmune myocarditis. [Clin Exp Pharmacol Physiol. 43 \(1\): 47-55.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10041 available at:  
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA1209F\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M368182:200529'

**Printed on 22 Mar 2021**