

Datasheet: MCA52GA

BATCH NUMBER 154015

| | |
|----------------------|---------------------|
| Description: | MOUSE ANTI RAT CD5 |
| Specificity: | CD5 |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | OX-19 |
| Isotype: | IgG1 |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | 1/100 |
| Immunohistology - Frozen | ▪ | | | |
| Immunohistology - Paraffin (1) | ▪ | | | |
| ELISA | | | ▪ | |
| Immunoprecipitation | ▪ | | | |
| Western Blotting | | | ▪ | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their particular system using appropriate negative/positive controls.

(1)The use of PLP fixation is recommended for optimal results.

| | |
|---------------------------------|---|
| Target Species | Rat |
| Product Form | Purified IgG - liquid |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A |
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | 0.09% Sodium Azide |

| | |
|---------------------------------------|--|
| Carrier Free | Yes |
| Approx. Protein Concentrations | IgG concentration 1.0mg/ml |
| Immunogen | Rat thymocyte glycoproteins. |
| External Database Links | <p>UniProt: P51882 Related reagents</p> <p>Entrez Gene: 54236 Cd5 Related reagents</p> |
| RRID | AB_566834 |
| Fusion Partners | Spleen cells from an immunized BALB/c mouse were fused with cells from the NS1 mouse myeloma cell line. |
| Specificity | <p>Mouse anti Rat CD5 antibody, clone OX-19 recognizes the rat CD5 cell surface antigen, a 69kD glycoprotein expressed by T cells, thymocytes and a subset of B cells.</p> <p>Mouse anti Rat CD5 antibody, clone OX-19 has been reported as being suitable for use on periodate-lysine paraformaldehyde (PLP) fixed paraffin embedded tissue (Whiteland et al. 1995).</p> <p>Mouse anti Rat CD5 antibody, clone OX-19 is routinely tested in flow cytometry on rat splenocytes.</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. |
| References | <ol style="list-style-type: none"> Dallman, M.J. <i>et al.</i> (1982) The roles of host and donor cells in the rejection of skin allografts by T cell-deprived rats injected with syngeneic T cells. Eur J Immunol. 12 (6): 511-8. Dallman, M.J. <i>et al.</i> (1984) MRC OX-19: a monoclonal antibody that labels rat T lymphocytes and augments in vitro proliferative responses. Eur J Immunol. 14 (3): 260-7. Huitinga, I. <i>et al.</i> (1995) Macrophages in T cell line-mediated, demyelinating, and chronic relapsing experimental autoimmune encephalomyelitis in Lewis rats. Clin Exp Immunol. 100 (2): 344-51. Whiteland, J.L. <i>et al.</i> (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. J Histochem Cytochem. 43 (3): 313-20. Song, E. <i>et al.</i> (2002) Early application of Met-RANTES ameliorates chronic allograft nephropathy. Kidney Int. 61: 676-85. Moreno-Manzano, V. <i>et al.</i> (2003) Retinoids as a potential treatment for experimental puromycin-induced nephrosis Br J Pharmacol. 139: 823-31. Shiraishi, H. <i>et al.</i> (2002) Antibody binding to fas ligand attenuates inflammatory cell infiltration and cytokine secretion, leading to reduction of myocardial infarct areas and |

- reperfusion injury. [Lab Invest. 82: 1121-9.](#)
8. Rusai, K. *et al.* (2008) Administration of interleukin-1 receptor antagonist ameliorates renal ischemia-reperfusion injury. [Transpl Int. 21: 572-80.](#)
9. Antus, B. *et al.* (2001) Contribution of androgens to chronic allograft nephropathy is mediated by dihydrotestosterone [Kidney Int. 60: 1955-63.](#)
10. Antus, B. *et al.* (2005) Effects of progesterone and selective oestrogen receptor modulators on chronic allograft nephropathy in rats. [Nephrol Dial Transplant. 20: 329-35.](#)
11. Kikuchi, H. *et al.* (2000) Severe proteinuria, sustained for 6 months, induces tubular epithelial cell injury and cell infiltration in rats but not progressive interstitial fibrosis. [Nephrol Dial Transplant. 15: 799-810.](#)
12. Abrams, M.B. *et al.* (2009) Multipotent mesenchymal stromal cells attenuate chronic inflammation and injury-induced sensitivity to mechanical stimuli in experimental spinal cord injury. [Restor Neurol Neurosci. 27: 307-21.](#)
13. Robichon, R. *et al.* (2005) Pig xenografts to the immunocompetent rat brain: Survival rates using distinct neurotoxic lesions in the nigrostriatal pathway and two rat strains. [Exp Neurol. 194: 333-40.](#)
14. de Vries, H.E. *et al.* (2002) Signal-regulatory protein alpha-CD47 interactions are required for the transmigration of monocytes across cerebral endothelium. [J Immunol. 168: 5832-9.](#)
15. Pamukcu, O. *et al.* (2016) Anti-inflammatory role of obestatin in autoimmune myocarditis. [Clin Exp Pharmacol Physiol. 43 \(1\): 47-55.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA52GA>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

- Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
- Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)
- Goat Anti Mouse IgG (STAR76...) [RPE](#)
- Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
- Goat Anti Mouse IgG (STAR70...) [FITC](#)

Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

| | | | | | |
|----------------------------------|---|------------------|---|---------------|---|
| North & South America | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com | Worldwide | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com | Europe | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com |
|----------------------------------|---|------------------|---|---------------|---|

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M368184:200529'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)