# Datasheet: MCA547B BATCH NUMBER 150907

Description:	MOUSE ANTI HUMAN CD34:Biotin
Specificity:	CD34 CLASS II
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	QBEND/10
Isotype:	lgG1
Quantity:	100 TESTS

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>							
	rad-antibodies.com/protocols.							
	Flass Ordensedure	Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	•		· · · · · · · · · · · · · · · · · · ·	Neat			
	Where this antibody has necessarily exclude its u			•	•			
	a guide only. It is recom		•		•			
	system using appropriate			-				
	system using appropriate	e negative/	positive					
Target Species	Human							
Species Cross	s Reacts with: Cynomolgus monkey, Rhesus Monkey							
Reactivity	Does not react with:Bovine, Sheep, Rat, Dog <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for							
	further information.							
Product Form	Purified IgG conjugated	to Biotin -	liquid					
Buffer Solution	Phosphate buffered salir	ne						
Preservative	0.09% Sodium Azide							
Stabilisers	1% Bovine Serum Albumin							
Approx. Protein	IgG concentration 0.1mg	ı/ml						

#### Concentrations

Immunogen	Human endothelial cell membrane vesicles.				
External Database Links	UniProt: <u>P28906</u> <u>Related reagents</u>				
	Entrez Gene: <u>947</u> CD34 <u>Related reagents</u>				
RRID	AB_2074373				
Fusion Partners	Spleen cells from immunized NZB mice were fused with cells of the mouse NSO myeloma cell line.				
Specificity	<b>Mouse anti Human CD34 antibody, clone QBEND/10</b> recognizes the human CD34 antigen, also known as Hematopoietic progenitor cell antigen CD34. Human CD34 is 385 amino acid polypeptide containing a 31 residue signal peptide, cleaved to yield the ~110kDa mature form of CD34, a sialomucin single pass transmembrane glycoprotein. CD34 is expressed by stem cells (Kaufman <i>et al.</i> 2001) and small vessel endothelium (Ramani <i>et al.</i> 1990)				
	Human CD34 exists as two isoforms, the full length form described here and a truncated isoform lacking the carboxy-terminal of the intracellular domain and containing some alternative sequence in the remaining intracellular region. Antibody binding epitopes on human CD34 have been classified according to their resistance to enzymatic degradation and grouped together using this and competitive binding assays (Lanza <i>et al.</i> 1999). Mouse anti Human CD34 antibody, clone QBEND/10 has been classified as binding to the class II epitope, resistant to neuraminidase treatment but sensitive to both glycoprotease and chymopapain digestion. Mouse anti Human CD34, clone 581 (MCA1578) which binds to the class III epitope resistant to all three enzymzatic treatments (Nishio <i>et al.</i> 1996 In Leukocyte Typing VI). Clone QBEND 10 is expected to bind to both isoforms of human CD34 as it's binding epitope has been mapped to the extracellular domain between amino acids 43 and 49 by peptide microarray analysis (Jones <i>et al.</i> 1996, in Leukocyte Typing VI).				
	Mouse anti Human CD34 antibody, clone QBEND/10 has been successfully exploited for the detection of CD34 in brain capillaries of Alzheimer's patients ( <u>Kalaria <i>et al.</i> 1992</u> ) and in acute lymphoblastic leukemia cells ( <u>Sutherland <i>et al.</i> 1992</u> ) by western blotting.				
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.				
References	<ol> <li>Fina, L. <i>et al.</i> (1990) Expression of the CD34 gene in vascular endothelial cells. <u>Blood.</u> <u>75 (12): 2417-26.</u></li> <li>Sopper, S. <i>et al.</i> (1997) Lymphocyte subsets and expression of differentiation markers in blood and lymphoid organs of rhesus monkeys. <u>Cytometry. 29 (4): 351-62.</u></li> </ol>				

3. Chan-Ling T (2011) Role of CD44+ Stem Cells in Mural Cell Formation in the Human Choroid: Evidence of Vascular Instability Due to Limited Pericyte Ensheathment. <u>Invest</u> <u>Ophthalmol Vis Sci. 52: 399-410.</u>

4. Pammer, J. *et al.* (1996) CD40 antigen is expressed by endothelial cells and tumor cells in Kaposi's sarcoma. <u>Am J Pathol. 148 (5): 1387-96.</u>

5. Lee, M.Y. et al. (2009) Angiogenesis in differentiated placental multipotent

mesenchymal stromal cells is dependent on integrin alpha5beta1. PLoS One. 4: e6913.

6. Chan-Ling, T. *et al.* (2004) Astrocyte-endothelial cell relationships during human retinal vascular development. <u>Invest Ophthalmol Vis Sci. 45: 2020-32.</u>

7. Chen, S.P. *et al.* (2014) Reduced circulating endothelial progenitor cells in reversible cerebral vasoconstriction syndrome. <u>J Headache Pain. 15: 82.</u>

8. Sauer, G. *et al.* (2003) Progression of cervical carcinomas is associated with down-regulation of CD9 but strong local re-expression at sites of transendothelial invasion. <u>Clin Cancer Res. 9: 6426-31.</u>

9. Sauter, B. *et al.* (1998) Immunoelectron Microscopic Characterization of Human Dermal Lymphatic Microvascular Endothelial Cells: Differential Expression of CD31, CD34, and Type IV Collagen with Lymphatic Endothelial Cells vs Blood Capillary Endothelial Cells in Normal Human Skin, Lymphangioma, and Hemangioma *In Situ*. J Histochem Cytochem. 46: 165-76.

10. Shetty, S. *et al.* (2011) Common lymphatic endothelial and vascular endothelial receptor-1 mediates the transmigration of regulatory T cells across human hepatic sinusoidal endothelium. J Immunol. 186: 4147-55.

11. Zhao, M. *et al.* (2007) Evidence for the presence of stem cell-like progenitor cells in human adult pancreas. <u>J Endocrinol. 195: 407-14.</u>

12. Jokubaitis, V.J. *et al.* (2008) Angiotensin-converting enzyme (CD143) marks hematopoietic stem cells in human embryonic, fetal, and adult hematopoietic tissues. <u>Blood. 111: 4055-63.</u>

13. Rutella, S. *et al.* (2003) Identification of a novel subpopulation of human cord blood CD34-CD133-CD7-CD45+lineage- cells capable of lymphoid/NK cell differentiation after in vitro exposure to IL-15. <u>J Immunol. 171: 2977-88.</u>

14. Suzuki, M. *et al.* (2012) Induction of human humoral immune responses in a novel HLA-DR-expressing transgenic NOD/Shi-scid/γcnull mouse. <u>Int Immunol. 24 (4): 243-52.</u>
15. Hsieh, J.Y. *et al.* (2013) miR-146a-5p circuitry uncouples cell proliferation and migration, but not differentiation, in human mesenchymal stem cells. <u>Nucleic Acids Res.</u> 41 (21): 9753-63.

16. Blank A et al. (2010) SDHB loss predicts malignancy in

pheochromocytomas/sympathethic paragangliomas, but not through hypoxia signalling. Endocr Relat Cancer. 17 (4): 919-28.

17. Junaid TO *et al.* (2014) Fetoplacental vascular alterations associated with fetal growth restriction. <u>Placenta. 35 (10): 808-15.</u>

18. Beleut M *et al.* (2012) Integrative genome-wide expression profiling identifies three distinct molecular subgroups of renal cell carcinoma with different patient outcome. <u>BMC</u> <u>Cancer. 12: 310.</u>

19. Chan-Ling T *et al.* (2011) Evidence of hematopoietic differentiation, vasculogenesis and angiogenesis in the formation of human choroidal blood vessels. <u>Exp Eye Res. 92 (5)</u>: 361-76.

20. Motamedian, S.R. et al. (2016) Response of Dental Pulp Stem Cells to Synthetic,

	Allograft, and Xenograft Bone Scaffolds. Int J Periodontics Restorative Dent. 37 (1): 49-59.
	21. Fan, C-Y. <i>et al.</i> (2017) <i>De novo</i> protein sequencing, humanization and <i>in vitro</i> effects
	of an antihuman CD34 mouse monoclonal antibody Biochemistry and Biophysics Reports.
	<u>9: 51-60.</u>
	22. Sameshima, N. et al. (2011) So-called 'adenosarcoma' of the kidney a novel adult
	renal tumor with a cystic appearance. Pathol Int. 61 (5): 313-8.
	23. Grognuz, A. et al. (2016) Human Fetal Progenitor Tenocytes for Regenerative
	Medicine. <u>Cell Transplant. 25 (3): 463-79.</u>
	24. Wang, D.Y. et al. (2017) Histological component quantification for the evaluation of
	endometrial receptivity in women with natural cycles undergoing in vitro
	fertilization/intracytoplasmic sperm injection. <u>Taiwan J Obstet Gynecol. 56 (3): 368-370.</u>
	25. GarikipatiV, N.S. <i>et al.</i> (2018) Isolation and characterization of mesenchymal stem
	cells from human fetus heart. <u>PLoS One. 13 (2): e0192244.</u>
	26. Rodewald, A.K. <i>et al.</i> (2019) Eight autopsy cases of melanoma brain metastases
	demonstrating angiotropism and pericytic mimicry. Implications for extravascular migratory
	metastasis. J Cutan Pathol. Mar 30 [Epub ahead of print].
Further Deeding	
Further Reading	1. Gorr, T.A. <i>et al.</i> (2011) Old proteins - new locations: myoglobin, haemoglobin, neuroglobin and cytoglobin in solid tumours and cancer cells. <u>Acta Physiol (Oxf). 202:</u> <u>563-581.</u>
Storage	neuroglobin and cytoglobin in solid tumours and cancer cells. Acta Physiol (Oxf). 202:
	neuroglobin and cytoglobin in solid tumours and cancer cells. <u>Acta Physiol (Oxf). 202:</u> <u>563-581.</u>
	neuroglobin and cytoglobin in solid tumours and cancer cells. <u>Acta Physiol (Oxf). 202:</u> 563-581. Store at +4°C or at -20°C if preferred.
	neuroglobin and cytoglobin in solid tumours and cancer cells. <u>Acta Physiol (Oxf). 202:</u> <u>563-581.</u> Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend
	neuroglobin and cytoglobin in solid tumours and cancer cells. <u>Acta Physiol (Oxf). 202:</u> <u>563-581.</u> Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may
	neuroglobin and cytoglobin in solid tumours and cancer cells. <u>Acta Physiol (Oxf). 202:</u> <u>563-581.</u> Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend
Storage	neuroglobin and cytoglobin in solid tumours and cancer cells. <u>Acta Physiol (Oxf). 202:</u> <u>563-581.</u> Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Storage Guarantee	neuroglobin and cytoglobin in solid tumours and cancer cells. <u>Acta Physiol (Oxf). 202:</u> 563-581. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch
Storage Guarantee Health And Safety	neuroglobin and cytoglobin in solid tumours and cancer cells. Acta Physiol (Oxf). 202: 563-581. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch Material Safety Datasheet documentation #10041 available at:
Storage Guarantee Health And Safety	neuroglobin and cytoglobin in solid tumours and cancer cells. <u>Acta Physiol (Oxf). 202:</u> 563-581. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch Material Safety Datasheet documentation #10041 available at: <u>https://www.bio-rad-antibodies.com/SDS/MCA547B</u>

## **Related Products**

### **Recommended Useful Reagents**

### HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rac	d.com	Email: antibody_sales_uk@bio-ra	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368201:200529'

#### Printed on 21 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint