

## Datasheet: MCA547F

**BATCH NUMBER 1711**

<b>Description:</b>	MOUSE ANTI HUMAN CD34:FITC
<b>Specificity:</b>	CD34 CLASS II
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	QBEND/10
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS/1ml

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

#### Target Species

Human

#### Species Cross Reactivity

Reacts with: Cynomolgus monkey, Rhesus Monkey

Does not react with: Bovine, Sheep, Rat, Dog

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

#### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

#### Buffer Solution

Phosphate buffered saline

<b>Preservative</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Stabilisers</b>	1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml
<b>Immunogen</b>	Human endothelial cell membrane vesicles.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P28906</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">947</a>    CD34    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_1125256
<b>Fusion Partners</b>	Spleen cells from immunized NZB mice were fused with cells of the mouse NSO myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human CD34 antibody, clone QBEND/10</b> recognizes the human CD34 antigen, also known as Hematopoietic progenitor cell antigen CD34. Human CD34 is 385 amino acid polypeptide containing a 31 residue signal peptide, cleaved to yield the ~110kDa mature form of CD34, a sialomucin single pass transmembrane glycoprotein. CD34 is expressed by stem cells (<a href="#">Kaufman <i>et al.</i> 2001</a>) and small vessel endothelium (<a href="#">Ramani <i>et al.</i> 1990</a>)</p> <p>Human CD34 exists as two isoforms, the full length form described here and a truncated isoform lacking the carboxy-terminal of the intracellular domain and containing some alternative sequence in the remaining intracellular region. Antibody binding epitopes on human CD34 have been classified according to their resistance to enzymatic degradation and grouped together using this and competitive binding assays (<a href="#">Lanza <i>et al.</i> 1999</a>). Mouse anti Human CD34 antibody, clone QBEND/10 has been classified as binding to the class II epitope, resistant to neuraminidase treatment but sensitive to both glycoprotease and chymopapain digestion. Mouse anti Human CD34 antibody, clone QBEND/10 binds to a different eoptope to Mouse anti Human CD34, clone 581 (<a href="#">MCA1578</a>) which binds to the class III epitope resistant to all three enzymzatic treatments (Nishio <i>et al.</i> 1996 In Leukocyte Typing VI). Clone QBEND 10 is expected to bind to both isoforms of human CD34 as it's binding epitope has been mapped to the extracellular domain between amino acids 43 and 49 by peptide microarray analysis (Jones <i>et al.</i> 1996, in Leukocyte Typing VI).</p> <p>Mouse anti Human CD34 antibody, clone QBEND/10 has been successfully exploited for the detection of CD34 in brain capillaries of Alzheimer's patients (<a href="#">Kalaria <i>et al.</i> 1992</a>) and in acute lymphoblastic leukemia cells (<a href="#">Sutherland <i>et al.</i> 1992</a>) by western blotting.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul

## References

1. Fina, L. *et al.* (1990) Expression of the CD34 gene in vascular endothelial cells. [Blood. 75 \(12\): 2417-26.](#)
2. Sopper, S. *et al.* (1997) Lymphocyte subsets and expression of differentiation markers in blood and lymphoid organs of rhesus monkeys. [Cytometry. 29 \(4\): 351-62.](#)
3. Chan-Ling T (2011) Role of CD44+ Stem Cells in Mural Cell Formation in the Human Choroid: Evidence of Vascular Instability Due to Limited Pericyte Ensheathment. [Invest Ophthalmol Vis Sci. 52: 399-410.](#)
4. Pammer, J. *et al.* (1996) CD40 antigen is expressed by endothelial cells and tumor cells in Kaposi's sarcoma. [Am J Pathol. 148 \(5\): 1387-96.](#)
5. Lee, M.Y. *et al.* (2009) Angiogenesis in differentiated placental multipotent mesenchymal stromal cells is dependent on integrin alpha5beta1. [PLoS One. 4: e6913.](#)
6. Chan-Ling, T. *et al.* (2004) Astrocyte-endothelial cell relationships during human retinal vascular development. [Invest Ophthalmol Vis Sci. 45: 2020-32.](#)
7. Chen, S.P. *et al.* (2014) Reduced circulating endothelial progenitor cells in reversible cerebral vasoconstriction syndrome. [J Headache Pain. 15: 82.](#)
8. Sauer, G. *et al.* (2003) Progression of cervical carcinomas is associated with down-regulation of CD9 but strong local re-expression at sites of transendothelial invasion. [Clin Cancer Res. 9: 6426-31.](#)
9. Sauter, B. *et al.* (1998) Immunoelectron Microscopic Characterization of Human Dermal Lymphatic Microvascular Endothelial Cells: Differential Expression of CD31, CD34, and Type IV Collagen with Lymphatic Endothelial Cells vs Blood Capillary Endothelial Cells in Normal Human Skin, Lymphangioma, and Hemangioma *In Situ*. [J Histochem Cytochem. 46: 165-76.](#)
10. Shetty, S. *et al.* (2011) Common lymphatic endothelial and vascular endothelial receptor-1 mediates the transmigration of regulatory T cells across human hepatic sinusoidal endothelium. [J Immunol. 186: 4147-55.](#)
11. Zhao, M. *et al.* (2007) Evidence for the presence of stem cell-like progenitor cells in human adult pancreas. [J Endocrinol. 195: 407-14.](#)
12. Jokubaitis, V.J. *et al.* (2008) Angiotensin-converting enzyme (CD143) marks hematopoietic stem cells in human embryonic, fetal, and adult hematopoietic tissues. [Blood. 111: 4055-63.](#)
13. Rutella, S. *et al.* (2003) Identification of a novel subpopulation of human cord blood CD34-CD133-CD7-CD45+lineage- cells capable of lymphoid/NK cell differentiation after in vitro exposure to IL-15. [J Immunol. 171: 2977-88.](#)
14. Suzuki, M. *et al.* (2012) Induction of human humoral immune responses in a novel HLA-DR-expressing transgenic NOD/Shi-scid/γnull mouse. [Int Immunol. 24 \(4\): 243-52.](#)
15. Hsieh, J.Y. *et al.* (2013) miR-146a-5p circuitry uncouples cell proliferation and migration, but not differentiation, in human mesenchymal stem cells. [Nucleic Acids Res. 41 \(21\): 9753-63.](#)
16. Blank A *et al.* (2010) SDHB loss predicts malignancy in pheochromocytomas/sympathetic paragangliomas, but not through hypoxia signalling. [Endocr Relat Cancer. 17 \(4\): 919-28.](#)
17. Junaid TO *et al.* (2014) Fetoplacental vascular alterations associated with fetal growth restriction. [Placenta. 35 \(10\): 808-15.](#)
18. Belet M *et al.* (2012) Integrative genome-wide expression profiling identifies three distinct molecular subgroups of renal cell carcinoma with different patient outcome. [BMC Cancer. 12: 310.](#)

19. Chan-Ling T *et al.* (2011) Evidence of hematopoietic differentiation, vasculogenesis and angiogenesis in the formation of human choroidal blood vessels. [Exp Eye Res. 92 \(5\): 361-76.](#)
20. Motamedian, S.R. *et al.* (2016) Response of Dental Pulp Stem Cells to Synthetic, Allograft, and Xenograft Bone Scaffolds. [Int J Periodontics Restorative Dent. 37 \(1\): 49-59.](#)
21. Fan, C-Y. *et al.* (2017) *De novo* protein sequencing, humanization and *in vitro* effects of an antihuman CD34 mouse monoclonal antibody [Biochemistry and Biophysics Reports. 9: 51-60.](#)
22. Sameshima, N. *et al.* (2011) So-called 'adenosarcoma' of the kidney a novel adult renal tumor with a cystic appearance. [Pathol Int. 61 \(5\): 313-8.](#)
23. Grognez, A. *et al.* (2016) Human Fetal Progenitor Tenocytes for Regenerative Medicine. [Cell Transplant. 25 \(3\): 463-79.](#)
24. Wang, D.Y. *et al.* (2017) Histological component quantification for the evaluation of endometrial receptivity in women with natural cycles undergoing *in vitro* fertilization/intracytoplasmic sperm injection. [Taiwan J Obstet Gynecol. 56 \(3\): 368-370.](#)
25. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. [PLoS One. 13 \(2\): e0192244.](#)
26. Rodewald, A.K. *et al.* (2019) Eight autopsy cases of melanoma brain metastases demonstrating angiotropism and pericytic mimicry. Implications for extravascular migratory metastasis. [J Cutan Pathol. Mar 30 \[Epub ahead of print\].](#)

---

**Further Reading** 1. Gorr, T.A. *et al.* (2011) Old proteins - new locations: myoglobin, haemoglobin, neuroglobin and cytoglobin in solid tumours and cancer cells. [Acta Physiol \(Oxf\). 202: 563-581.](#)

---

**Storage** Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA547F>  
10041

---

**Regulatory** For research purposes only

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

## Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M368202:200529'

Printed on 21 Feb 2024

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)