Datasheet: MCA547PET BATCH NUMBER 1704

Description:	MOUSE ANTI HUMAN CD34:RPE
Specificity:	CD34 CLASS II
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	QBEND/10
Isotype:	lgG1
Quantity:	25 TESTS/0.25ml

Product Details

Applications	s. This information is ions or personal dicated for further					
	information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry	-			Neat	
	Where this antibody has not been tested for use in a particular technique this does no necessarily exclude its use in such procedures. Suggested working dilutions are give a guide only. It is recommended that the user titrates the antibody for use in his/her o system using appropriate negative/positive controls.					
Target Species	Human					
Species Cross	Reacts with: Cynomol	gus monkey,	Rhesus N	lonkey		
ReactivityDoes not react with:Bovine, Sheep, Rat, DogN.B. Antibody reactivity and working conditions may vary between species. C						
				species. Cross		
	reactivity is derived from testing within our laboratories, peer-reviewed publications or					
	personal communications from the originators. Please refer to references indi				rences indicated for	
further information.						
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilised					
Reconstitution	Reconstitute with 0.25 ml distilled water					
Max Ex/Em	Fluorophore	Excitation M	ax (nm)	Emission Max (nm)		
	RPE 488nm laser	496		578		
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture					

	supernatant			
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	0.09% Sodium Azide1% Bovine Serum Albumin5% Sucrose			
Immunogen	Human endothelial cell membrane vesicles.			
External Database Links	UniProt: <u>P28906</u> <u>Related reagents</u> Entrez Gene: <u>947</u> CD34 <u>Related reagents</u>			
RRID	AB 2062999			
Fusion Partners	Spleen cells from immunized NZB mice were fused with cells of the mouse NSO myeloma cell line.			
Specificity	Mouse anti Human CD34 antibody, clone QBEND/10 recognizes the human CD34 antigen, also known as Hematopoietic progenitor cell antigen CD34. Human CD34 is 385 amino acid polypeptide containing a 31 residue signal peptide, cleaved to yield the ~110kDa mature form of CD34, a sialomucin single pass transmembrane glycoprotein. CD34 is expressed by stem cells (Kaufman <i>et al.</i> 2001) and small vessel endothelium (Ramani <i>et al.</i> 1990)			
	Human CD34 exists as two isoforms, the full length form described here and a truncated isoform lacking the carboxy-terminal of the intracellular domain and containing some alternative sequence in the remaining intracellular region. Antibody binding epitopes on human CD34 have been classified according to their resistance to enzymatic degradation and grouped together using this and competitive binding assays (Lanza <i>et al.</i> 1999). Mouse anti Human CD34 antibody, clone QBEND/10 has been classified as binding to the class II epitope, resistant to neuraminidase treatment but sensitive to both glycoprotease and chymopapain digestion. Mouse anti Human CD34, clone 581 (MCA1578) which binds to the class III epitope resistant to all three enzymzatic treatments (Nishio <i>et al.</i> 1996 In Leukocyte Typing VI). Clone QBEND 10 is expected to bind to both isoforms of human CD34 as it's binding epitope has been mapped to the extracellular domain between amino acids 43 and 49 by peptide microarray analysis (Jones <i>et al.</i> 1996, in Leukocyte Typing VI).			
	Mouse anti Human CD34 antibody, clone QBEND/10 has been successfully exploited for the detection of CD34 in brain capillaries of Alzheimer's patients (<u>Kalaria <i>et al.</i> 1992</u>) and			

Mouse anti Human CD34 antibody, clone QBEND/10 has been successfully exploited for the detection of CD34 in brain capillaries of Alzheimer's patients (<u>Kalaria *et al.* 1992</u>) and in acute lymphoblastic leukemia cells (<u>Sutherland *et al.* 1992</u>) by western blotting.

Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	1. Fina, L. et al. (1990) Expression of the CD34 gene in vascular endothelial cells. Blood
	<u>75 (12): 2417-26.</u>
	2. Sopper, S. et al. (1997) Lymphocyte subsets and expression of differentiation markers
	in blood and lymphoid organs of rhesus monkeys. Cytometry. 29 (4): 351-62.
	3. Chan-Ling T (2011) Role of CD44+ Stem Cells in Mural Cell Formation in the Human
	Choroid: Evidence of Vascular Instability Due to Limited Pericyte Ensheathment. Invest
	Ophthalmol Vis Sci. 52: 399-410.
	4. Pammer, J. et al. (1996) CD40 antigen is expressed by endothelial cells and tumor ce
	in Kaposi's sarcoma. <u>Am J Pathol. 148 (5): 1387-96.</u>
	5. Lee, M.Y. et al. (2009) Angiogenesis in differentiated placental multipotent
	mesenchymal stromal cells is dependent on integrin alpha5beta1. PLoS One. 4: e6913.
	6. Chan-Ling, T. et al. (2004) Astrocyte-endothelial cell relationships during human retina
	vascular development. Invest Ophthalmol Vis Sci. 45: 2020-32.
	7. Chen, S.P. et al. (2014) Reduced circulating endothelial progenitor cells in reversible
	cerebral vasoconstriction syndrome. J Headache Pain. 15: 82.
	8. Sauer, G. et al. (2003) Progression of cervical carcinomas is associated with
	down-regulation of CD9 but strong local re-expression at sites of transendothelial
	invasion. <u>Clin Cancer Res. 9: 6426-31.</u>
	9. Sauter, B. et al. (1998) Immunoelectron Microscopic Characterization of Human Derm
	Lymphatic Microvascular Endothelial Cells: Differential Expression of CD31, CD34, and
	Type IV Collagen with Lymphatic Endothelial Cells vs Blood Capillary Endothelial Cells in
	Normal Human Skin, Lymphangioma, and Hemangioma In Situ. J Histochem Cytochem.
	<u>46: 165-76.</u>
	10. Shetty, S. et al. (2011) Common lymphatic endothelial and vascular endothelial
	receptor-1 mediates the transmigration of regulatory T cells across human hepatic
	sinusoidal endothelium. <u>J Immunol. 186: 4147-55.</u>
	11. Zhao, M. et al. (2007) Evidence for the presence of stem cell-like progenitor cells in
	human adult pancreas. <u>J Endocrinol. 195: 407-14.</u>
	12. Jokubaitis, V.J. et al. (2008) Angiotensin-converting enzyme (CD143) marks
	hematopoietic stem cells in human embryonic, fetal, and adult hematopoietic tissues.
	Blood. 111: 4055-63.
	13. Rutella, S. <i>et al.</i> (2003) Identification of a novel subpopulation of human cord blood
	CD34-CD133-CD7-CD45+lineage- cells capable of lymphoid/NK cell differentiation after
	vitro exposure to IL-15. J Immunol. 171: 2977-88.
	14. Suzuki, M. <i>et al.</i> (2012) Induction of human humoral immune responses in a novel
	HLA-DR-expressing transgenic NOD/Shi-scid/ycnull mouse. Int Immunol. 24 (4): 243-52
	15. Hsieh, J.Y. <i>et al.</i> (2013) miR-146a-5p circuitry uncouples cell proliferation and
	migration, but not differentiation, in human mesenchymal stem cells. Nucleic Acids Res.
	41 (21): 9753-63.
	16. Blank A <i>et al.</i> (2010) SDHB loss predicts malignancy in
	pheochromocytomas/sympathethic paragangliomas, but not through hypoxia signalling.
	Endocr Relat Cancer. 17 (4): 919-28.
	17. Junaid TO <i>et al.</i> (2014) Fetoplacental vascular alterations associated with fetal growth
	restriction. <u>Placenta. 35 (10): 808-15.</u>
	18. Beleut M <i>et al.</i> (2012) Integrative genome-wide expression profiling identifies three
	15. Belout in et al. (2012) integrative generation entression promining identifies tillee

	 distinct molecular subgroups of renal cell carcinoma with different patient outcome. <u>BMC</u> <u>Cancer. 12: 310.</u> 19. Chan-Ling T <i>et al.</i> (2011) Evidence of hematopoietic differentiation, vasculogenesis and angiogenesis in the formation of human choroidal blood vessels. <u>Exp Eye Res. 92 (5): 361-76.</u> 20. Motamedian, S.R. <i>et al.</i> (2016) Response of Dental Pulp Stem Cells to Synthetic, Allograft, and Xenograft Bone Scaffolds. <u>Int J Periodontics Restorative Dent. 37 (1): 49-59.</u> 21. Fan, C-Y. <i>et al.</i> (2017) <i>De novo</i> protein sequencing, humanization and <i>in vitro</i> effects of an antihuman CD34 mouse monoclonal antibody <u>Biochemistry and Biophysics Reports.</u> 9: 51-60. 22. Sameshima, N. <i>et al.</i> (2011) So-called 'adenosarcoma' of the kidney a novel adult renal tumor with a cystic appearance. <u>Pathol Int. 61 (5): 313-8.</u> 23. Grognuz, A. <i>et al.</i> (2016) Human Fetal Progenitor Tenocytes for Regenerative Medicine. <u>Cell Transplant. 25 (3): 463-79.</u> 24. Wang, D.Y. <i>et al.</i> (2017) Histological component quantification for the evaluation of endometrial receptivity in women with natural cycles undergoing <i>in vitro</i> fertilization/intracytoplasmic sperm injection. <u>Taiwan J Obstet Gynecol. 56 (3): 368-370.</u> 25. GarikipatiV, N.S. <i>et al.</i> (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. <u>PLoS One. 13 (2): e0192244.</u> 26. Rodewald, A.K. <i>et al.</i> (2019) Eight autopsy cases of melanoma brain metastases demonstrating angiotropism and pericytic mimicry. Implications for extravascular migratory metastasis. J Cutan Pathol. Mar 30 [Epub ahead of print].
Further Reading	1. Gorr, T.A. <i>et al.</i> (2011) Old proteins - new locations: myoglobin, haemoglobin, neuroglobin and cytoglobin in solid tumours and cancer cells. <u>Acta Physiol (Oxf). 202:</u> <u>563-581.</u>
Storage	Prior to reconstitution store at +4°C. After reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA547PET 20487
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

Recommended Useful Reagents

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad	.com	Email: antibody_sales_uk@bio-rac	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375584:210104'

Printed on 21 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint