

Datasheet: MCA551T

Description:	MOUSE ANTI HUMAN CD11b
Specificity:	CD11b
Other names:	INTEGRIN ALPHA M CHAIN, MAC-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	ICRF44
Isotype:	IgG1
Quantity:	20 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/100
Immunohistology - Frozen (1)	▪			1/100 - 1/1000
Immunohistology - Paraffin		▪		

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Cynomolgus monkey, Baboon, Rhesus Monkey</p> <p>Does not react with: Cat</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Rheumatoid synovial cells and human monocytes
External Database Links	<p>UniProt: P11215 Related reagents</p> <p>Entrez Gene: 3684 ITGAM Related reagents</p>
Synonyms	CD11B, CR3A
RRID	AB_324415
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse Sp2/0 myeloma cell line
Specificity	<p>Mouse anti Human CD11b antibody, clone ICRF44 recognizes the human CD11b cell surface glycoprotein, a 165 kDa molecule also known as the alphaM integrin, MAC-1 and CR3. This molecule is expressed as a heterodimer in association with the beta 2 integrin, and is found upon monocytes, granulocytes, NK cells and some peripheral blood lymphocytes.</p> <p>Mouse anti Human CD11b antibody, clone ICRF44 has been reported to have various functional effects on monocytes, blocking adhesion and stimulating cytokine and chemokine release.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood
Histology Positive Control Tissue	Human tonsil
References	<ol style="list-style-type: none"> Yoshino, M. <i>et al.</i> (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (<i>Macaca fascicularis</i>) by using anti-human cross-reactive antibodies. Exp Anim. 49: 97-110. Jonker, M. <i>et al.</i> (1989) Reactivity of mAb specific for human CD markers with Rhesus monkey leucocyte. Leucocyte Typing IV. Oxford University Press 1058-1063.

3. Dransfield, I. *et al.* (1992) Interaction of leukocyte integrins with ligand is necessary but not sufficient for function. [J Cell Biol. 116 \(6\): 1527-35.](#)
4. Malhotra, V. *et al.* (1986) Ligand binding by the p150,95 antigen of U937 monocytic cells: properties in common with complement receptor type 3 (CR3). [Eur J Immunol. 16 \(9\): 1117-23.](#)
5. Glasow, A. *et al.* (2005) Retinoids and myelomonocytic growth factors co-operatively activate RAR{alpha} and induce human myeloid leukemia cell differentiation via MAP kinase pathways. [Blood 105: 341-9.](#)
6. Rezzonico, R. *et al.* (2001) Ligation of CD11b and CD11c beta(2) integrins by antibodies or soluble CD23 induces macrophage inflammatory protein 1alpha (MIP-1alpha) and MIP-1beta production in primary human monocytes through a pathway dependent on nuclear factor-kappaB. [Blood. 97 \(10\): 2932-40.](#)
7. Rezzonico, R. *et al.* (2000) Engagement of CD11b and CD11c beta2 integrin by antibodies or soluble CD23 induces IL-1beta production on primary human monocytes through mitogen-activated protein kinase-dependent pathways. [Blood. 95 \(12\): 3868-77.](#)
8. Canalli, A.A. *et al.* (2001) Participation of Mac-1, LFA-1 and VLA-4 integrins in the in vitro adhesion of sickle cell disease neutrophils to endothelial layers, and reversal of adhesion by simvastatin. [Haematologica 96: 526-33.](#)
9. Patel, S. *et al.* (2009) Reconstituted high-density lipoprotein increases plasma high-density lipoprotein anti-inflammatory properties and cholesterol efflux capacity in patients with type 2 diabetes. [J Am Coll Cardiol. 53: 962-71.](#)
10. Stirling, R.G. *et al.* (2001) Interleukin-5 induces CD34(+) eosinophil progenitor mobilization and eosinophil CCR3 expression in asthma. [Am J Respir Crit Care Med. 164: 1403-9.](#)
11. Urquhart, P. *et al.* (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. [J Pharmacol Exp Ther. 321: 656-62.](#)
12. Woollard, K.J. *et al.* (2002) Direct modulatory effect of C-reactive protein on primary human monocyte adhesion to human endothelial cells. [Clin Exp Immunol. 130: 256-62.](#)
13. Ramacciotti, E. *et al.* (2011) Evaluation of soluble p-selectin as a marker for the diagnosis of deep venous thrombosis. [Clin Appl Thromb Hemost. 17: 425-31.](#)
14. Paul, G. *et al.* (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. [PLoS One 7: e35577.](#)
15. Gomes-Alves, P. *et al.* (2016) *In vitro* expansion of human cardiac progenitor cells: Exploring 'omics tools for characterization of cell-based allogeneic products [Translational Research. 9 Feb \[Epub ahead of print\]](#)
16. Hughes, S.F. *et al.* (2020) The role of phagocytic leukocytes following flexible ureteroscopy, for the treatment of kidney stones: an observational, clinical pilots-study. [Eur J Med Res. 25 \(1\): 68.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®680](#),
[DyLight®800](#), [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M368227:200529'

Printed on 23 Mar 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)