

Datasheet: MCA554F BATCH NUMBER 1610

MOUSE ANTI HUMAN CD35:FITC		
CD35		
CR1		
FITC		
Monoclonal Antibody		
Monoclonal Antibody E11		
•		

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes No	Not Determined	Suggested Dilution	
	Flow Cytometry	-		Neat	
	Immunohistology - Froze	n			
	Immunohistology - Parafi	ïn			
Target Species	necessarily exclude its	use in such procedune in such procedu	ser titrates the antibody	g dilutions are given as	
Species Cross Reactivity	Reacts with: Cynomolgus monkey, Rhesus Monkey, Baboon N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.				
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	FITC	490	525		
Preparation	Purified IgG prepared	by affinity chromatog	raphy on Protein A fror	n tissue culture	

	supernatant			
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin			
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml			
Immunogen	Human acute monocytic leukaemia cells.			
External Database Links	UniProt: P17927 Related reagents Entrez Gene: 1378 CR1 Related reagents			
Synonyms	C3BR			
RRID	AB_321212			
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.			
Specificity	Mouse anti Human CD35 antibody, clone E11 recognizes complement receptor type 1, also known as CD35 or the C3b/C4b receptor. CD35 a single chain, type I single pass transmembrane glycoprotein which exists in four allotypic forms (A,B,C,D) of ~190 kDa, ~220 kDa, ~160 kDa and ~250 kDa respectively (Khera <i>et al.</i> 2009). CD35 is expressed by granulocytes, monocytes, B cells and some T cells. Functionally, CD35 acts as a receptor for the C3b component of complement.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.			
References	 Birmingham, D.J. <i>et al.</i> (1996) The baboon erythrocyte complement receptor is a glycophosphatidylinositol-linked protein encoded by a homologue of the human CR1-like genetic element. J Immunol. 157 (6): 2586-92. Sopper, S. <i>et al.</i> (1997) Lymphocyte subsets and expression of differentiation markers in blood and lymphoid organs of rhesus monkeys. Cytometry. 29 (4): 351-62. Weissgerber P <i>et al.</i> (2003) Investigation of mechanisms involved in phagocytosis of <i>Legionella pneumophila</i> by human cells. FEMS Microbiol Lett. 219 (2): 173-9. Schmidt, C.Q. <i>et al.</i> (2016) Selectivity of C3-opsonin targeted complement inhibitors: A distinct advantage in the protection of erythrocytes from paroxysmal nocturnal hemoglobinuria patients. Immunobiology. 221 (4): 503-11. Noda, G.S. <i>et al.</i> (2020) Specificities and isotypes of erythrocytes autoantibodies in patients with warm autoimmune hemolytic anemia Rev Cubana Hematol Inmunol Hemoter 			

	 <u>36(4): e1283</u>. Klaas, M. <i>et al.</i> (2023) Sialoadhesin (CD169/Siglec-1) is an elescapes inhibitory cis-interactions and synergizes with other mapromote phagocytosis. <u>Glycoconj J. 40 (2): 213-23.</u> Wagstaffe, H.R. <i>et al.</i> (2023) The Development of Immunologithe Level and Function of Antibodies Induced by <i>Klebsiella pner</i> Vaccines. <u>mSphere. : e0068022.</u> 	acrophage receptors to gical Assays to Evaluate	
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.		
Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA554F 10041		
Regulatory	For research purposes only		

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070	<u>A)</u>
HUMAN SEROBLOCK (BUF070	B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody_sales_uk@bio-ra	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M414905:221214'

Printed on 18 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint