

Datasheet: MCA55A488

Description:	MOUSE ANTI RAT CD4 (DOMAIN 1):Alexa Fluor® 488
Specificity:	CD4 (DOMAIN 1)
Format:	ALEXA FLUOR® 488
Product Type:	Monoclonal Antibody
Clone:	W3/25
Isotype:	lgG1
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat				
Product Form	Purified IgG conjugate	Purified IgG conjugated to Alexa Fluor® 488 - liquid			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nr	m)	
	Alexa Fluor®488	495	519		
Preparation	Purified IgG prepared supernatant	by affinity chromatog	raphy on Protein A	from tissue culture	
Buffer Solution	Phosphate buffered s	aline			
Preservative	0.09% Sodium Azide				
Stabilisers	1% Bovine Serum	Albumin			
Approx. Protein Concentrations	IgG concentration 0.0	5 mg/ml			

lm	m	ıın	OΠ	en
		ull	vу	CII

Rat Thymocyte Membrane Glycoproteins.

External Database Links

UniProt:

P05540 Related reagents

Entrez Gene:

24932 Cd4 Related reagents

RRID

AB 323825

Fusion Partners

Spleen cells from immunized BALB/c mouse were fused with cells of the mouse NS-1 myeloma cell line.

Specificity

Mouse anti Rat CD4 antibody, clone W3/25 recognizes the rat CD4 cell surface glycoprotein, a ~55 kDa molecule expressed by helper T cells and weakly by monocytes. This antibody inhibits proliferation and IL-2 production in the MLR reaction.

Mouse anti Rat CD4 antibody, clone W3/25 has been described reacting with paraffinembedded material following PLP fixation (periodate-lysine-paraformaldehyde) (Whiteland et al. 1995).

Mouse anti Rat CD4 antibody, clone W3/25 is routinely tested in flow cytometry on rat splenocytes.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

- 1. Williams, A.F. et al. (1977) Analysis of cell surfaces by xenogeneic myeloma-hybrid antibodies: differentiation antigens of rat lymphocytes. Cell. 12 (3): 663-73.
- 2. Barclay, A.N. (1981) The localization of populations of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues. Immunology. 42 (4): 593-600.
- 3. Pelegrí, C. et al. (1995) Immunohistochemical changes in synovial tissue during the course of adjuvant arthritis. J Rheumatol. 22 (1): 124-32.
- 4. Whiteland, J.L. et al. (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. J Histochem Cytochem. 43 (3): 313-20.
- 5. Bjersing, J.L. et al. (2002) Loss of ileal IgA+ plasma cells and of CD4+ lymphocytes in ileal Peyer's patches of vitamin A deficient rats. Clin Exp Immunol. 130 (3): 404-8.
- 6. Hofmann, N. et al. (2002) Increased expression of ICAM-1, VCAM-1, MCP-1, and MIP-1 alpha by spinal perivascular macrophages during experimental allergic encephalomyelitis in rats. BMC Immunol. 3: 11.
- 7. Banerjee, S. et al. (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. Br J Ophthalmol. 87 (12): 1515-22.
- 8. Ishizuka, S. et al. (2004) Fermentable dietary fiber potentiates the localization of immune cells in the rat large intestinal crypts. Exp Biol Med (Maywood). 229 (9): 876-84.
- 9. Latta, M. et al. (2007) CXCR6 is expressed on T cells in both T helper type 1 (Th1) inflammation and allergen-induced Th2 lung inflammation but is only a weak mediator of chemotaxis. Immunology. 121 (4): 555-64.

- 10. Bode, U. *et al.* (2008) Dendritic cell subsets in lymph nodes are characterized by the specific draining area and influence the phenotype and fate of primed T cells. Immunology. 123 (4): 480-90.
- 11. Zilka, N. *et al.* (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. <u>J</u> Neuroimmunol. 209 (1-2): 16-25.
- 12. Schwartzkopff, J. *et al.* (2010) NK cell depletion delays corneal allograft rejection in baby rats. Mol Vis. 16: 1928-35.
- 13. Schmiedl, A. *et al.* (2021) Lung development and immune status under chronic LPS exposure in rat pups with and without CD26/DPP4 deficiency. <u>Cell Tissue Res. Oct 04</u> [Epub ahead of print].
- 14. Matsuyama, S. *et al.* (2021) Properties of macrophages and lymphocytes appearing in rat renal fibrosis followed by repeated injection of cisplatin. J Vet Med Sci. 83 (9): 1435-42.
- 15. Cąkała-Jakimowicz, M. & Puzianowska-Kuznicka, M. (2022) Towards Understanding the Lymph Node Response to Skin Infection with Saprophytic *Staphylococcus epidermidis.*. Biomedicines. 10 (5): 1021.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee

12 months from date of despatch

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA55A488 10041

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL: Alexa Fluor® 488 (MCA1209A488)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 To

Email: antibody_sales_us@bio-rad.com Email: antibody_sales_uk@bio-rad.com Email: antibody_sales_de@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385898:210513'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint