

Datasheet: MCA55PB

BATCH NUMBER 0515

Description:	MOUSE ANTI RAT CD4 (DOMAIN 1):Pacific Blue®				
Specificity:	CD4 (DOMAIN 1)				
Format:	Pacific Blue®				
Product Type:	Monoclonal Antibody				
Clone:	W3/25				
Isotype:	IgG1				
Quantity:	100 TESTS/1ml				

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat				
Product Form	Purified IgG conjugated to Pacific Blue® - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm		
	Pacific Blue®	410	455		
reparation	Purified IgG prepare supernatant	d by affinity chromatog	raphy on Protein A t		
ffer Solution	Phosphate buffered	saline			
eservative	0.09% Sodium Azide	9			
tabilisers	1% Bovine Serun	n Albumin			
pprox. Protein oncentrations	IgG concentration 0.	05 mg/ml			

Immunoger	1
-----------	---

Rat Thymocyte Membrane Glycoproteins.

External Database

Links

UniProt:

P05540 Related reagents

Entrez Gene:

24932 Cd4 Related reagents

RRID

AB 567277

Fusion Partners

Spleen cells from immunized BALB/c mouse were fused with cells of the mouse NS-1 myeloma cell line.

Specificity

Mouse anti Rat CD4 antibody, clone W3/25 recognizes the rat CD4 cell surface glycoprotein, a ~55 kDa molecule expressed by helper T cells and weakly by monocytes. This antibody inhibits proliferation and IL-2 production in the MLR reaction.

Mouse anti Rat CD4 antibody, clone W3/25 has been described reacting with paraffinembedded material following PLP fixation (periodate-lysine-paraformaldehyde) (Whiteland et al. 1995).

Mouse anti Rat CD4 antibody, clone W3/25 is routinely tested in flow cytometry on rat splenocytes.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

- 1. Williams, A.F. *et al.* (1977) Analysis of cell surfaces by xenogeneic myeloma-hybrid antibodies: differentiation antigens of rat lymphocytes. Cell. 12 (3): 663-73.
- 2. Barclay, A.N. (1981) The localization of populations of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues. Immunology. 42 (4): 593-600.
- 3. Whiteland, J.L. *et al.* (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. <u>J Histochem Cytochem. 43 (3): 313-20.</u>
- 4. Pelegrí, C. *et al.* (1995) Immunohistochemical changes in synovial tissue during the course of adjuvant arthritis. J Rheumatol. 22 (1): 124-32.
- 5. Hofmann, N. *et al.* (2002) Increased expression of ICAM-1, VCAM-1, MCP-1, and MIP-1 alpha by spinal perivascular macrophages during experimental allergic encephalomyelitis in rats. <u>BMC Immunol. 3: 11.</u>
- 6. Zilka, N. *et al.* (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. <u>J</u> Neuroimmunol. 209 (1-2): 16-25.
- 7. Schwartzkopff, J. *et al.* (2010) NK cell depletion delays corneal allograft rejection in baby rats. Mol Vis. 16: 1928-35.
- 8. Banerjee, S. *et al.* (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. <u>Br J Ophthalmol. 87 (12): 1515-22.</u>
- 9. Bjersing, J.L. *et al.* (2002) Loss of ileal IgA+ plasma cells and of CD4+ lymphocytes in ileal Peyer's patches of vitamin A deficient rats. <u>Clin Exp Immunol</u>. 130 (3): 404-8.

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:

https://www.bio-rad-antibodies.com/SDS/MCA55PB

10041

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:Pacific Blue® (MCA1209PB)

America

North & South Tel: +1 800 265 7376 Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368253:200529'

Printed on 18 Jan 2024