

## Datasheet: MCA55XZ

<b>Description:</b>	MOUSE ANTI RAT CD4 (DOMAIN 1):Preservative Free
<b>Specificity:</b>	CD4 (DOMAIN 1)
<b>Format:</b>	Preservative Free
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	W3/25
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	1 mg

## Product Details

**RRID** AB\_321378

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			
Functional Assays	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)PLP fixation is recommended for optimal results, see [Whiteland et al.](#) for details

**Target Species** Rat

**Product Form** Purified IgG - liquid

**Preparation** Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

**Buffer Solution** Phosphate buffered saline

**Preservative Stabilisers** None present

**Carrier Free** Yes

**Approx. Protein Concentrations** IgG concentration 1.0 mg/ml

<b>Immunogen</b>	Rat Thymocyte Membrane Glycoproteins.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P05540</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">24932</a>    Cd4    <a href="#">Related reagents</a></p>
<b>Fusion Partners</b>	Spleen cells from immunized BALB/c mouse were fused with cells of the mouse NS-1 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Rat CD4 antibody, clone W3/25</b> recognizes the rat CD4 cell surface glycoprotein, a ~55 kDa molecule expressed by helper T cells and weakly by monocytes. This antibody inhibits proliferation and IL-2 production in the MLR reaction.</p> <p>Mouse anti Rat CD4 antibody, clone W3/25 has been described reacting with paraffin-embedded material following PLP fixation (periodate-lysine-paraformaldehyde) (<a href="#">Whiteland et al. 1995</a>).</p> <p>Mouse anti Rat CD4 antibody, clone W3/25 is routinely tested in flow cytometry on rat splenocytes.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>Williams, A.F. <i>et al.</i> (1977) Analysis of cell surfaces by xenogeneic myeloma-hybrid antibodies: differentiation antigens of rat lymphocytes. <a href="#">Cell. 12 (3): 663-73.</a></li> <li>Barclay, A.N. (1981) The localization of populations of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues. <a href="#">Immunology. 42 (4): 593-600.</a></li> <li>Whiteland, J.L. <i>et al.</i> (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. <a href="#">J Histochem Cytochem. 43 (3): 313-20.</a></li> <li>Pelegrí, C. <i>et al.</i> (1995) Immunohistochemical changes in synovial tissue during the course of adjuvant arthritis. <a href="#">J Rheumatol. 22 (1): 124-32.</a></li> <li>Hofmann, N. <i>et al.</i> (2002) Increased expression of ICAM-1, VCAM-1, MCP-1, and MIP-1 alpha by spinal perivascular macrophages during experimental allergic encephalomyelitis in rats. <a href="#">BMC Immunol. 3: 11.</a></li> <li>Zilka, N. <i>et al.</i> (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. <a href="#">J Neuroimmunol. 209 (1-2): 16-25.</a></li> <li>Schwartzkopff, J. <i>et al.</i> (2010) NK cell depletion delays corneal allograft rejection in baby rats. <a href="#">Mol Vis. 16: 1928-35.</a></li> <li>Banerjee, S. <i>et al.</i> (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. <a href="#">Br J Ophthalmol. 87 (12): 1515-22.</a></li> <li>Bjersing, J.L. <i>et al.</i> (2002) Loss of ileal IgA+ plasma cells and of CD4+ lymphocytes in ileal Peyer's patches of vitamin A deficient rats. <a href="#">Clin Exp Immunol. 130 (3): 404-8.</a></li> </ol>
<b>Storage</b>	<p>Store at -20°C only.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>

<b>Guarantee</b>	18 months from date of despatch.
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10162 available at: 10162: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf</a>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

- Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
- Goat Anti Mouse IgG (STAR77...) [HRP](#)
- Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
- Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)
- Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
- Goat Anti Mouse IgG (STAR76...) [RPE](#)
- Goat Anti Mouse IgG (STAR70...) [FITC](#)
- Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
- Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
- Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®680](#),  
[DyLight®800](#), [FITC](#), [HRP](#)

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Preservative Free \(MCA1209XZ\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M341719:190109'

Printed on 20 May 2019