

Datasheet: MCA5646PE

Description:	RAT ANTI MOUSE CD276:RPE	
Specificity:	CD276	
Other names:	B7-H3	
Format:	RPE	
Product Type:	Monoclonal Antibody	
Clone:	MJ8	
Isotype:	lgG2b	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Links

UniProt:

Q8VE98 Related reagents

Entrez Gene:

102657 Cd276 Related reagents

Synonyms	B7h3
RRID	AB_10961292
Fusion Partners	Lymph node cells from immunised Sprague Dawley rats were fused with cells of the P3U1 myeloma cell line.
Specificity	Rat anti Mouse CD276 antibody, clone MJ8 recognizes mouse CD276, otherwise known as B7-H3, a ubiquitously expressed transmembrane glycoprotein and member of the B7 family of co-stimulatory molecules, which acts as both a positive and negative regulator of T-cell-mediated immune responses.
	CD276 is highly expressed in bone during embryogenesis, and can be induced on dendritic cells and monocytes by inflammatory cytokines. CD276 has been implicated in the development of acute and chronic transplant rejection, and is reported to have therapeutic potential as a regulator of cell-mediated immune responses to cancer, particularly in conjunction with anti-angiogenic therapy. In mice, CD276 has been linked with the development of pathogenic Th2 cells during the induction phase of allergic asthma (Nagashima et al. 2008).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
	The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A or BUF041B).
References	1. Nagashima, O. <i>et al.</i> (2008) B7-H3 contributes to the development of pathogenic Th2 cells in a murine model of asthma. <u>J Immunol. 181 (6): 4062-71.</u>
Storage	Prior to reconstitution store at +4°C. After reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A)
MOUSE SEROBLOCK FcR (BUF041B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 'M368275:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | Legal | Imprint