

## Datasheet: MCA5646PE

Description:	RAT ANTI MOUSE CD276:RPE
Specificity:	CD276
Other names:	B7-H3
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	MJ8
Isotype:	lgG2b
Quantity:	100 TESTS

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	•						
	Where this product has			•	•			
	necessarily exclude its a guide only. It is recor system using appropria	mmended the	at the use	er titrates the product f	g dilutions are given as or use in their own			
Target Species	Mouse							
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - Iyophilized							
Reconstitution	Reconstitute with 1ml distilled water							
Max Ex/Em	Fluorophore	Excitation M	lax (nm)	Emission Max (nm)				
	RPE 488nm laser	496		578				
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant							
Buffer Solution	Phosphate buffered sa	line						
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum A 5% Sucrose	Albumin						

Immunogen	Mouse IgG2a Fc-CD276 (aa 1-242).				
External Database Links	UniProt:				
LIIKJ	Q8VE98 Related reagents				
	Entrez Gene:				
	<u>102657</u> Cd276 <u>Related reagents</u>				
Synonyms	B7h3				
RRID	AB_10961292				
Fusion Partners	Lymph node cells from immunised Sprague Dawley rats were fused with cells of the P3U1 myeloma cell line.				
Specificity	<b>Rat anti Mouse CD276 antibody, clone MJ8</b> recognizes mouse CD276, otherwise known as B7-H3, a ubiquitously expressed transmembrane glycoprotein and member of the B7 family of co-stimulatory molecules, which acts as both a positive and negative regulator of T-cell-mediated immune responses.				
	CD276 is highly expressed in bone during embryogenesis, and can be induced on dendritic cells and monocytes by inflammatory cytokines. CD276 has been implicated in the development of acute and chronic transplant rejection, and is reported to have therapeutic potential as a regulator of cell-mediated immune responses to cancer, particularly in conjunction with anti-angiogenic therapy. In mice, CD276 has been linked with the development of pathogenic Th2 cells during the induction phase of allergic asthma (Nagashima <i>et al.</i> 2008).				
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.				
	The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR ( <u>BUF041A</u> or <u>BUF041B</u> ).				
References	1. Nagashima, O. <i>et al</i> . (2008) B7-H3 contributes to the development of pathogenic Th2 cells in a murine model of asthma. <u>J Immunol. 181 (6): 4062-71.</u>				
Storage	Prior to reconstitution store at +4°C. After reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.				
Guarantee	12 months from date of despatch				
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA5646PE">https://www.bio-rad-antibodies.com/SDS/MCA5646PE</a>				

20487						
For rese	For research purposes only					
Products						
ended Useful Rea	gents					
ROBLOCK FcR (BUFC	<u>)41A)</u>					
ROBLOCK FcR (BUF(	<u>)41B)</u>					
el: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21		
ax: +1 919 878 3751 mail: antibody_sales_us@bi		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50		
	Products ended Useful Rea ROBLOCK FcR (BUFC ROBLOCK FcR (BUFC	Products ended Useful Reagents ROBLOCK FcR (BUF041A) ROBLOCK FcR (BUF041B)	Products   ended Useful Reagents   ROBLOCK FcR (BUF041A)   ROBLOCK FcR (BUF041B)   ei: +1 800 265 7376   Worldwide   Tel: +44 (0)1865 852 700	Products   ended Useful Reagents   ROBLOCK FcR (BUF041A)   ROBLOCK FcR (BUF041B)   al: +1 800 265 7376   Worldwide   Tel: +44 (0)1865 852 700   Europe		

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint