

Datasheet: MCA5656

Description:	MOUSE ANTI BOVINE MHC CLASS II DR
Specificity:	MHC CLASS II DR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CC108
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/200
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Functional Assays			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Bovine
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes

Approx. Protein Concentrations	IgG concentration 1.0mg/ml
External Database Links	UniProt: Q30309 Related reagents P79464 Related reagents
RRID	AB_10843456
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the Mouse NS1 myeloma cell line.
Specificity	<p>Mouse anti Bovine MHC class II DR antibody, clone CC108 recognizes Bovine MHC Class II DR. MHC Class II molecules are constitutively expressed on antigen presenting cells such as dendritic cells, B lymphocytes, monocytes, macrophages, activated T lymphocytes and may be induced on a range of other cell types by interferon gamma.</p> <p>The major histocompatibility complex (MHC) is a cluster of genes some of which are important in the immune response to infections. In cattle, this complex is referred to as the bovine leukocyte antigen (BoLA) region. There are 2 major types of MHC class IIa molecules encoded by the BoLA which are DR and DQ each composed of an alpha and beta chain.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	<ol style="list-style-type: none"> Stephens, S.A. & Howard, C.J. (2002) Infection and transformation of dendritic cells from bovine afferent lymph by <i>Theileria annulata</i>. Parasitology. 124 (Pt 5): 485-93. Yamakawa, Y. <i>et al.</i> (2008) Identification and functional characterization of a bovine orthologue to DC-SIGN. J Leukoc Biol. 83 (6): 1396-403. Corripio-Miyar, Y. <i>et al.</i> (2015) Phenotypic and functional analysis of monocyte populations in cattle peripheral blood identifies a subset with high endocytic and allogeneic T-cell stimulatory capacity. Vet Res. 46: 112. Guzman, E. <i>et al.</i> (2014) Bovine $\gamma\delta$ T cells are a major regulatory T cell subset. J Immunol. 193 (1): 208-22. Childerstone, A.J. <i>et al.</i> (1999) Demonstration of bovine CD8+ T-cell responses to foot-and-mouth disease virus. J Gen Virol. 80 (Pt 3): 663-9. Sopp, P. <i>et al.</i> (1994) Detection of bovine viral diarrhoea virus p80 protein in subpopulations of bovine leukocytes. J Gen Virol. 75 (Pt 5): 1189-94. Bembridge, G.P. <i>et al.</i> (1995) CD45RO expression on bovine T cells: relation to biological function. Immunology. 86 (4): 537-44. Gibson, A.J. <i>et al.</i> (2016) Differential macrophage function in Brown Swiss and Holstein Friesian cattle. Vet Immunol Immunopathol. 181: 15-23. Corripio-miyar, Y. <i>et al.</i> (2017) 1,25-Dihydroxyvitamin D3 modulates the phenotype and function of Monocyte derived dendritic cells in cattle BMC Veterinary Research. 13 (1) [Epub ahead of print]. Risalde, M.A. <i>et al.</i> (2020) BVDV permissiveness and lack of expression of co-stimulatory molecules on PBMCs from calves pre-infected with BVDV. Comp Immunol

[Microbiol Infect Dis. 68: 101388.](#)

11. Park, D.S. *et al.* (2021) Dynamic changes in blood immune cell composition and function in Holstein and Jersey steers in response to heat stress. [Cell Stress Chaperones. 26 \(4\): 705-20.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M384313:210513'

Printed on 08 Nov 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)