



Datasheet: MCA5665SBUV700

BATCH NUMBER 64673304

Description:	MOUSE ANTI HUMAN CD16:StarBright UltraViolet 700
Specificity:	CD16
Other names:	FcRIII
Format:	StarBright UltraViolet 700
Product Type:	Monoclonal Antibody
Clone:	KD1
Isotype:	IgG2a
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Bovine, Sheep, Dolphin Does not react with: Rat, Dog Reacts weakly with: Horse, Pig N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.		
Product Form	Purified IgG conjugated to StarBright UltraViolet 700 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	StarBright UltraViolet 700	340	700

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin 0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20
Approx. Protein Concentrations	For information on the concentration of our StarBright Dye conjugated reagents please visit our FAQ page.
Immunogen	A polyclonal population of NK cells.
External Database Links	UniProt: P79107 Related reagents P08637 Related reagents O75015 Related reagents Entrez Gene: 281766 FCGR3A Related reagents 2214 FCGR3A Related reagents 2215 FCGR3B Related reagents
Synonyms	CD16A, CD16B, FCG3, FCGR3, FCGRIII, IGFR3
Fusion Partners	Spleen cells from immunised mice were fused with cells of the P3U1 myeloma cell line.
Specificity	Mouse anti Human CD16 antibody, clone KD1 recognizes human CD16, a 50-65 kDa cell surface molecule, which is the low affinity receptor for IgG (FcR III). CD16 exists as a transmembranous form (Fc gammaRIIIA, or CD16A) and a glycosyl phosphatidylinositol (GPI) anchored form (Fc gammaRIIIB, or CD16B). CD16A is expressed by NK cells, some T cells, and macrophages (Moretta et al. 1990), whereas CD16B is primarily expressed by granulocytes (Bonecchi et al. 1999). Clone KD1 recognizes both forms of CD16 and will therefore recognize all cell types expressing CD16. Mouse anti Human CD16, clone KD1 can be used to identify CD16 in a range of species including bovine (Boysen et al. 2010) and ovine (Elhmouzi-Younes et al. 2010).
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	- Moretta, A. <i>et al.</i> (1989) CD16 surface molecules regulate the cytolytic function of CD3CD16+ human natural killer cells. Int J Cancer. 44 (4): 727-30. - Ciccone, E. <i>et al.</i> (1990) Specific recognition of human CD3-CD16+ natural killer cells

- requires the expression of an autosomic recessive gene on target cells. [J Exp Med. 172 \(1\): 47-52.](#)
3. Zocchi, M.R. *et al.* (1998) HIV-1 Tat inhibits human natural killer cell function by blocking L-type calcium channels. [J Immunol. 161: 2938-43.](#)
 4. Hernández-Caselles, T. *et al.* (2006) A study of CD33 (SIGLEC-3) antigen expression and function on activated human T and NK cells: two isoforms of CD33 are generated by alternative splicing. [J Leukoc Biol. 79: 46-58.](#)
 5. Boysen, P. *et al.* (2008) Natural killer cells in lymph nodes of healthy calves express CD16 and show both cytotoxic and cytokine-producing properties. [Dev Comp Immunol. 32: 773-83.](#)
 6. Connelley, T. *et al.* (2011) NKp46 defines ovine cells that have characteristics corresponding to NK cells. [Vet Res. 42: 37.](#)
 7. Gibson, A.J. *et al.* (2016) Differential macrophage function in Brown Swiss and Holstein Friesian cattle. [Vet Immunol Immunopathol. 181: 15-23.](#)
 8. Pomeroy, B. *et al.* (2016) Longitudinal characterization of bovine monocyte-derived dendritic cells from mid-gestation into subsequent lactation reveals nadir in phenotypic maturation and macrophage-like cytokine profile in late gestation. [J Reprod Immunol. 118: 1-8.](#)
 9. Elnaggar, M.M. *et al.* (2017) Identification of monoclonal antibodies cross-reactive with bottlenose dolphin orthologues of the major histocompatibility complex and leukocyte differentiation molecules. [Vet Immunol Immunopathol. 192: 54-9.](#)
 10. Pomeroy, B. *et al.* (2017) Counts of bovine monocyte subsets prior to calving are predictive for postpartum occurrence of mastitis and metritis. [Vet Res. 48 \(1\): 13.](#)
 11. Elnaggar, M.M. *et al.* (2019) Pattern of CD14, CD16, CD163 and CD172a expression on water buffalo (*Bubalus bubalis*.) leukocytes. [Vet Immunol Immunopathol. 211: 1-5.](#)
 12. Liu, J. *et al.* (2020) *Theileria annulata*. transformation altered cell surface molecules expression and endocytic function of monocyte-derived dendritic cells. [Ticks Tick Borne Dis. 11 \(3\): 101365.](#)
 13. Kolar, Q.K. *et al.* (2020) Anatomical distribution of respiratory tract leukocyte cell subsets in neonatal calves. [Vet Immunol Immunopathol. 227: 110090.](#)
 14. Park, D.S. *et al.* (2021) Dynamic changes in blood immune cell composition and function in Holstein and Jersey steers in response to heat stress. [Cell Stress Chaperones. 26 \(4\): 705-20.](#)
 15. Fiorenza, M.F. *et al.* (2021) Neutrophils recognize and amplify IFNT signals derived from day 7 bovine embryo for stimulation of ISGs expression *in vitro*.: A possible implication for the early maternal recognition of pregnancy. [Biochem Biophys Res Commun. 553: 37-43.](#)
 16. Howell, A. *et al.* (2024) Resting and activated bovine neutrophils and eosinophils differ in their responses to adrenergic agonists. [Vet Immunol Immunopathol. 272: 110758.](#)
 17. Hong, S. *et al.* (2024) Impact of an Injectable Trace Mineral Supplement on the Immune Response and Outcome of *Mannheimia haemolytica* Infection in Feedlot Cattle. [Biol Trace Elem Res. Jun 10 \[Epub ahead of print\].](#)
 18. Scatà, C.M. *et al.* (2024) Characterization of Cellular Immune System at Different Ages in Water Buffalo (*Bubalus bubalis*) [Journal of Buffalo Science. 13: 133-139.](#)
 19. Gilbert, F.B. *et al.* (2025) Expression of FcμR by bovine mononuclear blood leukocytes. [Dev Comp Immunol. 162: 105304.](#)
-

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA5665SBUV700
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M445019:250815'

Printed on 10 Jul 2026