Datasheet: MCA5672 BATCH NUMBER 152352

Description:	MOUSE ANTI HUMAN CD63
Specificity:	CD63
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	TEA3/18
lsotype:	lgG1
Quantity:	0.2 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is
	derived from testing within our laboratories, peer-reviewed publications or personal
	communications from the originators. Please refer to references indicated for further
	information. For general protocol recommendations, please visit <u>www.bio-</u>
	rad-antibodies.com/protocols.
	Yes No Not Determined Suggested Dilution

		res	NO	Not Determined	Suggested Dilution
	Flow Cytometry	•			Neat - 1/10
	Immunohistology - Frozen			•	
	Immunohistology - Paraffin			•	
	ELISA			•	
	Immunoprecipitation			•	
	Western Blotting				
	Where this product has r	not been t	ested for	use in a particular tec	hnique this does not
	necessarily exclude its u	se in such	n procedu	res. Suggested working	ng dilutions are given as
	a guide only. It is recomm	nended th	at the us	er titrates the product	for use in their own
	system using appropriate	e negative	/positive	controls.	
Target Species	Human				
Product Form	Purified IgG - liquid				
Buffer Solution	Phosphate buffered salin	e			
Preservative	<0.1% Sodium Azide (Na	aN ₃)			
Stabilisers	1% Bovine Serum Album	nin			
Approx. Protein Concentrations	IgG concentration 0.1mg	/ml			
Immunogen	TNF activated HUVEC co	ells			

External Database Links	UniProt:
	P08962 Related reagents
	Entrez Gene:
	<u>967</u> CD63 <u>Related reagents</u>
Synonyms	MLA1, TSPAN30
RRID	AB_10709460
Specificity	Mouse anti Human CD63 antibody, clone TEA3/18 recognizes the CD63 cell surface antigen, a glycoprotein of ~40-60 kDa.
	CD63 is expressed by monocytes and by a variety of cell lines. It is located intracellularly in lysosomal granules of platelets, being translocated to the surface upon activation. It may therefore by useful as a marker of platelet activation.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul or 100ul whole blood.
References	 Peñas, P.F. <i>et al.</i> (2000) Tetraspanins are localized at motility-related structures and involved in normal human keratinocyte wound healing migration. <u>J Invest Dermatol. 114</u> (6): 1126-35. Casey, T.M. <i>et al.</i> (2007) Organelle proteomics: identification of the exocytic machinery associated with the natural killer cell secretory lysosome. <u>Mol Cell Proteomics. 6 (5)</u>: <u>767-80.</u> Cepeda, V. & Fraile-Ramos, A. (2011) A role for the SNARE protein syntaxin 3 in human cytomegalovirus morphogenesis. <u>Cell Microbiol. 13 (6)</u>: 846-58. Suárez, H. <i>et al.</i> (2017) A bead-assisted flow cytometry method for the semi-quantitative analysis of Extracellular Vesicles. <u>Sci Rep. 7 (1)</u>: 11271. Oliveira-rodríguez, M. <i>et al.</i> (2016) Development of a rapid lateral flow immunoassay test for detection of exosomes previously enriched from cell culture medium and body fluids. <u>J Extracell Vesicles. 5: 31803.</u> Andreu, Z. <i>et al.</i> (2016) Comparative analysis of EV isolation procedures for miRNAs detection in serum samples. <u>J Extracell Vesicles. 5: 31655.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in
	frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA5672

	10041				
Regulato	ry For resea	irch purpose	s only		
Relate	d Products				
Recomm	nended Secondary A	Antibodies			
Rabbit A	nti Mouse IgG (STAR12) <u>RI</u>	<u>PE</u>		
Goat Ant	i Mouse IgG IgA IgM (S	TAR87) <u>H</u>	<u>RP</u>		
Goat Ant	i Mouse IgG (STAR76	.) <u>RI</u>	<u>PE</u>		
Rabbit A	nti Mouse IgG (STAR13	5) <u>H</u>	<u>RP</u>		
Goat Ant	i Mouse IgG (STAR70	.) <u>FI</u>	<u>TC</u>		
Goat Ant	i Mouse IgG (H/L) (STA		<u>k. Phos., DyLight®488, [</u> /Light®650, DyLight®68 TC, HRP		
Rabbit A	nti Mouse IgG (STAR9	.) FI	TC		
Goat Ant	i Mouse IgG (STAR77	.) <u>H</u>	<u>RP</u>		
Goat Ant	i Mouse IgG (Fc) (STAF	R120) <u>FI</u>	TC, <u>HRP</u>		
Recom	nended Negative Co	ntrols			
MOUSE Io	G1 NEGATIVE CONTRO	L (MCA928)			
orth & South merica	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-i	Worldwide rad.com	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.co
To find a b	atch/lot specific datashee	t for this prod	uct, please use our online s	search tool at:	bio-rad-antibodies.com/datashee

'M396053:220608'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint