Datasheet: MCA5672 BATCH NUMBER 159057

| Description:MOUSE ANTI HUMAN OSpecificity:CD63Format:PurifiedProduct Type:Monoclonal AntibodyClone:TEA3/18 | |
|--|------|
| Format:PurifiedProduct Type:Monoclonal Antibody | ;D63 |
| Product Type: Monoclonal Antibody | |
| | |
| Clone: TEA3/18 | |
| | |
| Isotype: IgG1 | |
| Quantity: 0.2 mg | |

Product Details

| Applications | This product has been reported to work in the following applications. This information is |
|--------------|---|
| | derived from testing within our laboratories, peer-reviewed publications or personal |
| | communications from the originators. Please refer to references indicated for further |
| | information. For general protocol recommendations, please visit <u>www.bio-</u> |
| | rad-antibodies.com/protocols. |
| | Yes No Not Determined Suggested Dilution |

| | | Yes | No | Not Determine | ed Suggested Dilution |
|-----------------------------------|----------------------------|-------------------|-----------|---------------------|-------------------------------|
| | Flow Cytometry | - | | | Neat - 1/10 |
| | Immunohistology - Frozen | | | • | |
| | Immunohistology - Paraffin | | | • | |
| | ELISA | | | • | |
| | Immunoprecipitation | | | • | |
| | Western Blotting | | | • | |
| | Where this product has r | not been te | ested for | use in a particula | r technique this does not |
| | necessarily exclude its u | se in such | n procedu | res. Suggested w | orking dilutions are given as |
| | a guide only. It is recomm | nended th | at the us | er titrates the pro | duct for use in their own |
| | system using appropriate | e negative | /positive | controls. | |
| Target Species | Human | | | | |
| Product Form | Purified IgG - liquid | | | | |
| Buffer Solution | Phosphate buffered salin | е | | | |
| Preservative | <0.1% Sodium Azide (Na | aN ₃) | | | |
| Stabilisers | 1% Bovine Serum Album | nin | | | |
| Approx. Protein Concentrations | IgG concentration 0.1mg | /ml | | | |
| Immunogen | TNF activated HUVEC c | ells | | | |

| External Database Links | UniProt: |
|----------------------------------|--|
| | P08962 Related reagents |
| | Entrez Gene: |
| | <u>967</u> CD63 <u>Related reagents</u> |
| Synonyms | MLA1, TSPAN30 |
| RRID | AB_10709460 |
| Specificity | Mouse anti Human CD63 antibody, clone TEA3/18 recognizes the CD63 cell surface antigen, a glycoprotein of ~40-60 kDa. |
| | CD63 is expressed by monocytes and by a variety of cell lines. It is located intracellularly in lysosomal granules of platelets, being translocated to the surface upon activation. It may therefore by useful as a marker of platelet activation. |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul or 100ul whole blood. |
| References | Peñas, P.F. <i>et al.</i> (2000) Tetraspanins are localized at motility-related structures and involved in normal human keratinocyte wound healing migration. <u>J Invest Dermatol. 114</u> (6): 1126-35. Casey, T.M. <i>et al.</i> (2007) Organelle proteomics: identification of the exocytic machinery associated with the natural killer cell secretory lysosome. <u>Mol Cell Proteomics. 6 (5)</u>: <u>767-80.</u> Cepeda, V. & Fraile-Ramos, A. (2011) A role for the SNARE protein syntaxin 3 in human cytomegalovirus morphogenesis. <u>Cell Microbiol. 13 (6)</u>: 846-58. Suárez, H. <i>et al.</i> (2017) A bead-assisted flow cytometry method for the semi-quantitative analysis of Extracellular Vesicles. <u>Sci Rep. 7 (1)</u>: 11271. Oliveira-rodríguez, M. <i>et al.</i> (2016) Development of a rapid lateral flow immunoassay test for detection of exosomes previously enriched from cell culture medium and body fluids. <u>J Extracell Vesicles. 5: 31803.</u> Andreu, Z. <i>et al.</i> (2016) Comparative analysis of EV isolation procedures for miRNAs detection in serum samples. <u>J Extracell Vesicles. 5: 31655.</u> |
| Storage | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in |
| | frost-free freezers is not recommended. |
| Guarantee | 12 months from date of despatch |
| Health And Safety Information | Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA5672 |

| | 10041 | | | | |
|------------------------|--|----------------------|--|-----------------|---|
| Regulato | ry For resea | irch purpose | | | |
| Relate | d Products | | | | |
| Recomm | nended Secondary A | Antibodies | | | |
| Rabbit A | nti Mouse IgG (STAR12 |) <u>RI</u> | <u>PE</u> | | |
| Goat Ant | i Mouse IgG IgA IgM (S | TAR87) <u>H</u> | <u>RP</u> | | |
| Goat Ant | i Mouse IgG (STAR76 | .) <u>RI</u> | <u>PE</u> | | |
| Rabbit A | nti Mouse IgG (STAR13 | 5) <u>H</u> | <u>RP</u> | | |
| Goat Ant | i Mouse IgG (STAR70 | .) <u>FI</u> | <u>TC</u> | | |
| Goat Ant | i Mouse IgG (H/L) (STA | | <u>k. Phos., DyLight®488, [</u> /Light®650, DyLight®68 TC, HRP | | |
| Rabbit A | nti Mouse IgG (STAR9 | .) FI | TC | | |
| Goat Ant | i Mouse IgG (STAR77 | .) <u>H</u> | <u>RP</u> | | |
| Goat Ant | i Mouse IgG (Fc) (STAF | R120) <u>FI</u> | TC, <u>HRP</u> | | |
| Recom | nended Negative Co | ntrols | | | |
| MOUSE Io | G1 NEGATIVE CONTRO | L (MCA928) | | | |
| orth & South merica | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-i | Worldwide rad.com | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio | Europe | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.co |
| To find a b | atch/lot specific datashee | t for this prod | uct, please use our online s | search tool at: | bio-rad-antibodies.com/datashee |

'M396053:220608'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint