



## Datasheet: MCA5705PE

**BATCH NUMBER 163063**

<b>Description:</b>	HAMSTER ANTI MOUSE DELTA-LIKE PROTEIN 1:RPE
<b>Specificity:</b>	DELTA-LIKE PROTEIN 1
<b>Other names:</b>	DLL1
<b>Format:</b>	RPE
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	HMD1-5
<b>Isotype:</b>	IgG
<b>Quantity:</b>	100 TESTS

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

### Target Species

Mouse

### Species Cross Reactivity

Reacts with: Rat

Reacts weakly with: Human

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

### Reconstitution

Reconstitute with 1.0ml distilled water

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578

**Preparation** Purified IgG prepared by affinity chromatography on Protein G

---

**Buffer Solution** Phosphate buffered saline

---

**Preservative** 0.09% Sodium Azide (NaN<sub>3</sub>)  
**Stabilisers** 1% Bovine Serum Albumin  
5% Sucrose

---

**Immunogen** DLL1-expressing CHO cells.

---

**External Database Links**

**UniProt:**

[Q61483](#) [Related reagents](#)

[P97677](#) [Related reagents](#)

[O00548](#) [Related reagents](#)

**Entrez Gene:**

[13388](#) DII1 [Related reagents](#)

[84010](#) DII1 [Related reagents](#)

[28514](#) DLL1 [Related reagents](#)

---

**RRID** AB\_11152602

---

**Fusion Partners** Spleen cells from immunised Armenian hamsters were fused with cells of the P3U1 myeloma cell line.

---

**Specificity** **Hamster anti Mouse Delta-Like Protein 1 antibody, clone HMD1-5** recognizes Delta-like protein 1 (DLL1), one of the five major ligands of the Notch signaling pathway, which is activated through the binding of specific ligands to the Notch receptors Notch 1-4.

The Notch signaling pathway is an evolutionarily conserved pathway in multi-cellular organisms, which is vital for cell-cell communication, important during fundamental developmental and physiological processes, including regulation of cell fate decisions during neuronal, cardiac and endocrine development, stem cell hematopoiesis, thymic T-cell development, and both tumor progression and suppression.

Ligation of Notch receptors by their specific ligands, Jagged1 (CD339), Jagged2, Delta-like protein 1 (DLL1), DLL3 and DLL4, on physically adjacent signal receiving cells, induces proteolysis of the receptors by ADAM-family metalloproteases and the gamma-secretase complex, within the transmembrane domain, releasing the Notch intracellular domain (NICD) to translocate to the nucleus. Subsequent signal transduction then occurs through either the CSL-NICD-Mastermind complex cascade (canonical pathway), or NF-kappaB-NICD and CSL-NICD-Deltex complex signaling cascades (non-canonical pathway). The canonical pathway inhibits the differentiation of stem cells or progenitor cells, whilst the non-canonical pathway promotes differentiation.

DLL1 is widely expressed, and acts as a mediator of cell fate decisions during hematopoiesis, and may play a role in cell-to-cell communication in mammalian embryos.

DLL1 plays an important role in B and T cell differentiation, in embryonic somite formation and patterning, and associates with the scaffolding protein MAGI1 at adherens junctions on neuronal processes. Signaling through DLL1 and Notch 2 has been implicated in the development of marginal zone B cells (MZB).

Hamster anti Mouse Delta-Like Protein 1 antibody, clone HMD1-5 blocks binding of Notch2 to Dll1 ([Moriyama et al. 2008](#))

---

**Flow Cytometry** Use 10ul of the suggested working dilution to label  $1 \times 10^6$  cells in 100ul.

---

**References**

1. Moriyama, Y. *et al.* (2008) Delta-like 1 is essential for the maintenance of marginal zone B cells in normal mice but not in autoimmune mice. [Int Immunol. 20 \(6\): 763-73.](#)
2. Sekine, C. *et al.* (2009) Differential regulation of splenic CD8- dendritic cells and marginal zone B cells by Notch ligands. [Int Immunol. 21 \(3\): 295-301.](#)
3. Sekine, C. *et al.* (2012) Differential regulation of osteoclastogenesis by Notch2/Delta-like 1 and Notch1/Jagged1 axes. [Arthritis Res Ther. 14: R45.](#)

---

**Further Reading**

1. Bray, S.J. (2006) Notch signalling: a simple pathway becomes complex. [Nat Rev Mol Cell Biol. 7 \(9\): 678-89.](#)
2. Iso, T. *et al.* (2003) Notch signaling in vascular development. [Arterioscler Thromb Vasc Biol. 23 \(4\): 543-53.](#)
3. Hu, X. *et al.* (2008) Integrated regulation of Toll-like receptor responses by Notch and interferon-gamma pathways. [Immunity. 29 \(5\): 691-703.](#)
4. Hoyne, G.F. *et al.* (2001) Notch signalling in the regulation of peripheral immunity. [Immunol Rev. 182: 215-27.](#)

---

**Storage** Prior to reconstitution store at +4°C.  
After reconstitution store at +4°C.  
DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #20487 available at: <https://www.bio-rad-antibodies.com/SDS/MCA5705PE>  
20487

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL:RPE \(MCA2356PE\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M375598:210104'

**Printed on 18 Jan 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)