

Datasheet: MCA5707GA

Description:	HAMSTER ANTI MOUSE CD339
Specificity:	CD339
Other names:	JAGGED1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	HMJ1-29
Isotype:	IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Mouse

Species Cross Reactivity

Reacts with: Rat

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein G

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Jagged1-expressing CHO cells.
External Database Links	<p>UniProt:</p> <p>Q9QXX0 Related reagents</p> <p>Q63722 Related reagents</p> <p>Entrez Gene:</p> <p>16449 Jag1 Related reagents</p> <p>29146 Jag1 Related reagents</p>
RRID	AB_10709279
Fusion Partners	Spleen cells from immunized Armenian hamsters were fused with cells of the P3U1 myeloma cell line.
Specificity	<p>Hamster anti Mouse CD339 antibody, clone HMJ1-29 specifically recognizes CD339, otherwise known as Jagged1, one of the five major ligands of the Notch signaling pathway, which is activated through the binding of specific ligands to the Notch receptors Notch 1-4.</p> <p>In humans, mutations in the JAG1 gene are responsible for the autosomal dominant multisystem disorder Alagille syndrome 1 (ALGS1)</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Mouse spleen
References	<ol style="list-style-type: none"> Moriyama Y <i>et al.</i> (2008) Delta-like 1 is essential for the maintenance of marginal zone B cells in normal mice but not in autoimmune mice. Int Immunol. 20 (6): 763-73. Sekine, C. <i>et al.</i> (2009) Differential regulation of splenic CD8- dendritic cells and marginal zone B cells by Notch ligands. Int Immunol. 21 (3): 295-301. Sekine, C. <i>et al.</i> (2012) Differential regulation of osteoclastogenesis by Notch2/Delta-like 1 and Notch1/Jagged1 axes. Arthritis Res Ther. 14: R45. Koyanagi, A. <i>et al.</i> (2012) Expression of Notch receptors and ligands on immature and mature T cells. Biochem Biophys Res Commun. 418: 799-805.
Further Reading	<ol style="list-style-type: none"> Hoyne, G.F. <i>et al.</i> (2001) Notch signalling in the regulation of peripheral immunity. Immunol Rev. 182: 215-27. Iso, T. <i>et al.</i> (2003) Notch signaling in vascular development. Arterioscler Thromb Vasc

[Biol. 23 \(4\): 543-53.](#)

3. Bray, S.J. (2006) Notch signalling: a simple pathway becomes complex. [Nat Rev Mol Cell Biol. 7 \(9\): 678-89.](#)

4. Hu, X. *et al.* (2008) Integrated regulation of Toll-like receptor responses by Notch and interferon-gamma pathways. [Immunity. 29 \(5\): 691-703.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA5707GA>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight@550](#), [DyLight@650](#), [DyLight@800](#),
[FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M384303:210513'

Printed on 05 Sep 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)