

Datasheet: MCA5708GA

BATCH NUMBER 170329

Description:	HAMSTER ANTI MOUSE JAGGED2
Specificity:	JAGGED2
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	HMJ2-1
Isotype:	IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Species Cross Reactivity	Reacts with: Rat, Human N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Jagged2-expressing CHO cells.
External Database Links	UniProt: Q9QYE5 Related reagents P97607 Related reagents Q9Y219 Related reagents Entrez Gene: 16450 Jag2 Related reagents 29147 Jag2 Related reagents 3714 JAG2 Related reagents
RRID	AB_10708971
Fusion Partners	Spleen cells from immunised Armenian hamsters were fused with cells of the P3U1 myeloma cell line.
Specificity	Hamster anti Mouse JAGGED2 antibody, clone HMJ2-1 recognizes Jagged2, one of the five major ligands of the Notch signaling pathway, which is activated through the binding of specific ligands to the Notch receptors Notch 1-4. The Notch signaling pathway is an evolutionarily conserved pathway in multi-cellular organisms, which is vital for cell-cell communication, important during fundamental developmental and physiological processes, including regulation of cell fate decisions during neuronal, cardiac and endocrine development, stem cell hematopoiesis, thymic T-cell development, and both tumor progression and suppression. Ligation of Notch receptors by their specific ligands, Jagged1 (CD339), Jagged2, Delta-like protein 1 (DLL1), DLL3 and DLL4, on physically adjacent signal receiving cells, induces proteolysis of the receptors by ADAM-family metalloproteases and the gamma-secretase complex, within the transmembrane domain, releasing the Notch intracellular domain (NICD) to translocate to the nucleus. Subsequent signal transduction then occurs through either the CSL-NICD-Mastermind complex cascade (canonical pathway), or NF-kappaB-NICD and CSL-NICD-Delta complex signaling cascades (non-canonical pathway). The canonical pathway inhibits the differentiation of stem cells or progenitor cells, whilst the non-canonical pathway promotes differentiation. Jagged2 is expressed by stromal and thymic lymphoid cells, and by splenic macrophages and dendritic cells (DCs), and plays a vital role during limb, craniofacial, tooth, and thymic

development, as well as being implicated in the maintenance and function of neuronal cells in both the central (CNS) and enteric (gastrointestinal) nervous system.

Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
Histology Positive Control Tissue	Mouse spleen
References	1. Moriyama, Y. <i>et al.</i> (2008) Delta-like 1 is essential for the maintenance of marginal zone B cells in normal mice but not in autoimmune mice. Int Immunol. 20 (6): 763-73. 2. Sekine, C. <i>et al.</i> (2009) Differential regulation of splenic CD8- dendritic cells and marginal zone B cells by Notch ligands. Int Immunol. 21 (3): 295-301. 3. Sekine, C. <i>et al.</i> (2012) Differential regulation of osteoclastogenesis by Notch2/Delta-like 1 and Notch1/Jagged1 axes. Arthritis Res Ther. 14: R45.
Further Reading	1. Sander, G.R. <i>et al.</i> (2003) Expression of Notch1 and Jagged2 in the enteric nervous system. J Histochem Cytochem. 51 (7): 969-72. 2. Bray, S.J. (2006) Notch signalling: a simple pathway becomes complex. Nat Rev Mol Cell Biol. 7 (9): 678-89. 3. Iso, T. <i>et al.</i> (2003) Notch signaling in vascular development. Arterioscler Thromb Vasc Biol. 23 (4): 543-53. 4. Hu, X. <i>et al.</i> (2008) Integrated regulation of Toll-like receptor responses by Notch and interferon-gamma pathways. Immunity. 29: 691-703 5. Hoyne, G.F. <i>et al.</i> (2001) Notch signalling in the regulation of peripheral immunity. Immunol Rev. 182: 215-27.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA5708GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight®550](#), [DyLight®650](#), [DyLight®800](#),
[FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M368335:200529'

Printed on 28 Aug 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)