

# Datasheet: MCA5708GA

**BATCH NUMBER 171003**

<b>Description:</b>	HAMSTER ANTI MOUSE JAGGED2
<b>Specificity:</b>	JAGGED2
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	HMJ2-1
<b>Isotype:</b>	IgG
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Species Cross Reactivity</b>	<p>Reacts with: Rat, Human</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
<b>Immunogen</b>	Jagged2-expressing CHO cells.
<b>External Database Links</b>	<p><b>UniProt:</b></p> <p><a href="#">Q9QYE5</a>   <a href="#">Related reagents</a></p> <p><a href="#">P97607</a>   <a href="#">Related reagents</a></p> <p><a href="#">Q9Y219</a>   <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b></p> <p><a href="#">16450</a>   Jag2   <a href="#">Related reagents</a></p> <p><a href="#">29147</a>   Jag2   <a href="#">Related reagents</a></p> <p><a href="#">3714</a>   JAG2   <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_10708971
<b>Fusion Partners</b>	Spleen cells from immunised Armenian hamsters were fused with cells of the P3U1 myeloma cell line.
<b>Specificity</b>	<p><b>Hamster anti Mouse JAGGED2 antibody, clone HMJ2-1</b> recognizes Jagged2, one of the five major ligands of the Notch signaling pathway, which is activated through the binding of specific ligands to the Notch receptors Notch 1-4.</p> <p>The Notch signaling pathway is an evolutionarily conserved pathway in multi-cellular organisms, which is vital for cell-cell communication, important during fundamental developmental and physiological processes, including regulation of cell fate decisions during neuronal, cardiac and endocrine development, stem cell hematopoiesis, thymic T-cell development, and both tumor progression and suppression.</p> <p>Ligation of Notch receptors by their specific ligands, Jagged1 (CD339), Jagged2, Delta-like protein 1 (DLL1), DLL3 and DLL4, on physically adjacent signal receiving cells, induces proteolysis of the receptors by ADAM-family metalloproteases and the gamma-secretase complex, within the transmembrane domain, releasing the Notch intracellular domain (NICD) to translocate to the nucleus. Subsequent signal transduction then occurs through either the CSL-NICD-Mastermind complex cascade (canonical pathway), or NF-kappaB-NICD and CSL-NICD-Delta complex signaling cascades (non-canonical pathway). The canonical pathway inhibits the differentiation of stem cells or progenitor cells, whilst the non-canonical pathway promotes differentiation.</p> <p>Jagged2 is expressed by stromal and thymic lymphoid cells, and by splenic macrophages and dendritic cells (DCs), and plays a vital role during limb, craniofacial, tooth, and thymic</p>

development, as well as being implicated in the maintenance and function of neuronal cells in both the central (CNS) and enteric (gastrointestinal) nervous system.

<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul.
<b>Histology Positive Control Tissue</b>	Mouse spleen
<b>References</b>	<ol style="list-style-type: none"><li>1. Moriyama, Y. <i>et al.</i> (2008) Delta-like 1 is essential for the maintenance of marginal zone B cells in normal mice but not in autoimmune mice. <a href="#">Int Immunol. 20 (6): 763-73.</a></li><li>2. Sekine, C. <i>et al.</i> (2009) Differential regulation of splenic CD8- dendritic cells and marginal zone B cells by Notch ligands. <a href="#">Int Immunol. 21 (3): 295-301.</a></li><li>3. Sekine, C. <i>et al.</i> (2012) Differential regulation of osteoclastogenesis by Notch2/Delta-like 1 and Notch1/Jagged1 axes. <a href="#">Arthritis Res Ther. 14: R45.</a></li></ol>
<b>Further Reading</b>	<ol style="list-style-type: none"><li>1. Sander, G.R. <i>et al.</i> (2003) Expression of Notch1 and Jagged2 in the enteric nervous system. <a href="#">J Histochem Cytochem. 51 (7): 969-72.</a></li><li>2. Bray, S.J. (2006) Notch signalling: a simple pathway becomes complex. <a href="#">Nat Rev Mol Cell Biol. 7 (9): 678-89.</a></li><li>3. Iso, T. <i>et al.</i> (2003) Notch signaling in vascular development. <a href="#">Arterioscler Thromb Vasc Biol. 23 (4): 543-53.</a></li><li>4. Hu, X. <i>et al.</i> (2008) Integrated regulation of Toll-like receptor responses by Notch and interferon-gamma pathways. <a href="#">Immunity. 29: 691-703</a></li><li>5. Hoyne, G.F. <i>et al.</i> (2001) Notch signalling in the regulation of peripheral immunity. <a href="#">Immunol Rev. 182: 215-27.</a></li></ol>
<b>Storage</b>	<p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p> <p>This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA5708GA">https://www.bio-rad-antibodies.com/SDS/MCA5708GA</a> 10040
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight®550](#), [DyLight®650](#), [DyLight®800](#),  
[FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

### Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M368335:200529'

**Printed on 28 Aug 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)