

## Datasheet: MCA5708GA

<b>Description:</b>	HAMSTER ANTI MOUSE JAGGED2
<b>Specificity:</b>	JAGGED2
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	HMJ2-1
<b>Isotype:</b>	IgG
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Species Cross Reactivity</b>	Reacts with: Rat, Human <b>N.B.</b> Antibody reactivity and working conditions may vary between species.
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
<b>Immunogen</b>	Jagged2-expressing CHO cells.

**External Database  
Links**

**UniProt:**

[Q9QYE5](#)    [Related reagents](#)  
[P97607](#)    [Related reagents](#)  
[Q9Y219](#)    [Related reagents](#)

**Entrez Gene:**

[16450](#)    Jag2    [Related reagents](#)  
[29147](#)    Jag2    [Related reagents](#)  
[3714](#)    JAG2    [Related reagents](#)

---

**RRID**

AB\_10708971

---

**Fusion Partners**

Spleen cells from immunised Armenian hamsters were fused with cells of the P3U1 myeloma cell line.

---

**Specificity**

**Hamster anti Mouse JAGGED2 antibody, clone HMJ2-1** recognizes Jagged2, one of the five major ligands of the Notch signaling pathway, which is activated through the binding of specific ligands to the Notch receptors Notch 1-4.

The Notch signaling pathway is an evolutionarily conserved pathway in multi-cellular organisms, which is vital for cell-cell communication, important during fundamental developmental and physiological processes, including regulation of cell fate decisions during neuronal, cardiac and endocrine development, stem cell hematopoiesis, thymic T-cell development, and both tumor progression and suppression.

Ligation of Notch receptors by their specific ligands, Jagged1 (CD339), Jagged2, Delta-like protein 1 (DLL1), DLL3 and DLL4, on physically adjacent signal receiving cells, induces proteolysis of the receptors by ADAM-family metalloproteases and the gamma-secretase complex, within the transmembrane domain, releasing the Notch intracellular domain (NICD) to translocate to the nucleus. Subsequent signal transduction then occurs through either the CSL-NICD-Mastermind complex cascade (canonical pathway), or NF-kappaB-NICD and CSL-NICD-Deltex complex signaling cascades (non-canonical pathway). The canonical pathway inhibits the differentiation of stem cells or progenitor cells, whilst the non-canonical pathway promotes differentiation.

Jagged2 is expressed by stromal and thymic lymphoid cells, and by splenic macrophages and dendritic cells (DCs), and plays a vital role during limb, craniofacial, tooth, and thymic development, as well as being implicated in the maintenance and function of neuronal cells in both the central (CNS) and enteric (gastrointestinal) nervous system.

---

**Flow Cytometry**

Use 10ul of the suggested working dilution to label  $1 \times 10^6$  cells in 100ul.

---

**Histology Positive  
Control Tissue**

Mouse spleen

---

**References**

1. Moriyama, Y. *et al.* (2008) Delta-like 1 is essential for the maintenance of marginal zone B cells in normal mice but not in autoimmune mice. [Int Immunol. 20 \(6\): 763-73.](#)
2. Sekine, C. *et al.* (2009) Differential regulation of splenic CD8- dendritic cells and marginal zone B cells by Notch ligands. [Int Immunol. 21 \(3\): 295-301.](#)
3. Sekine, C. *et al.* (2012) Differential regulation of osteoclastogenesis by Notch2/Delta-like 1 and Notch1/Jagged1 axes. [Arthritis Res Ther. 14: R45.](#)

**Further Reading**

1. Sander, G.R. *et al.* (2003) Expression of Notch1 and Jagged2 in the enteric nervous system. [J Histochem Cytochem. 51 \(7\): 969-72.](#)
2. Bray, S.J. (2006) Notch signalling: a simple pathway becomes complex. [Nat Rev Mol Cell Biol. 7 \(9\): 678-89.](#)
3. Iso, T. *et al.* (2003) Notch signaling in vascular development. [Arterioscler Thromb Vasc Biol. 23 \(4\): 543-53.](#)
4. Hu, X. *et al.* (2008) Integrated regulation of Toll-like receptor responses by Notch and interferon-gamma pathways. [Immunity. 29: 691-703](#)
5. Hoyne, G.F. *et al.* (2001) Notch signalling in the regulation of peripheral immunity. [Immunol Rev. 182: 215-27.](#)

---

**Storage** Store at +4°C or at -20°C if preferred.  
Storage in frost-free freezers is not recommended.  
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory** For research purposes only

## Related Products

### Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight®800](#), [FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

### Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M368335:200529'

**Printed on 11 Aug 2020**

---

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)