Datasheet: MCA5750Z BATCH NUMBER 169166

Description:	MOUSE ANTI HUMAN VEGF: Preservative Free		
Specificity:	VEGF		
Other names:	VPF		
Format:	Preservative Free		
Product Type:	Monoclonal Antibody		
Clone:	4B2-8		
Isotype:	lgG1		
Quantity:	0.5 mg		

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA				2.0 - 4.0ug/ml
Immunoprecipitation			•	
Western Blotting				0.25 - 0.50ug/ml
Functional Assays	-			0.1 - 0.2ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - lyophilized
Reconstitution	Reconstitute with 0.5 ml distilled water
	Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture

	supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	None present		
Carrier Free	Yes		
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml after reconstitution		
Immunogen	Highly pure recombinant human VEGF ₁₆₅ (PHP091).		
External Database Links	UniProt: P15692 Related reagents Entrez Gene: 7422 VEGFA Related reagents		
Synonyms	VEGF		
RRID	AB_10897188		
	Mouse anti Human Vascular endothelial growth factor antibody, clone 4B2-8 recognizes human Vascular endothelial growth factor A , also known as VEGF-A or Vascular permeability factor. VEGF-A is a 206 amino acid ~45 kDa disulphide linked homodimeric heparin-binding cytokine and member of the PDGF/VEGF growth factor family, which exists as several splice variants of the VEGF gene, and is a potent stimulator of angiogenesis of both normal and cancerous cells, and acts as a regulator of vasculogenesis.		
Specificity	recognizes human Vascular endothelial growth factor A , also known as VEGF-A or Vascular permeability factor. VEGF-A is a 206 amino acid ~45 kDa disulphide linked homodimeric heparin-binding cytokine and member of the PDGF/VEGF growth factor family, which exists as several splice variants of the VEGF gene, and is a potent stimulator of angiogenesis of both normal and cancerous cells, and acts as a regulator of		
Specificity	recognizes human Vascular endothelial growth factor A , also known as VEGF-A or Vascular permeability factor. VEGF-A is a 206 amino acid ~45 kDa disulphide linked homodimeric heparin-binding cytokine and member of the PDGF/VEGF growth factor family, which exists as several splice variants of the VEGF gene, and is a potent stimulator of angiogenesis of both normal and cancerous cells, and acts as a regulator of		
Specificity	recognizes human Vascular endothelial growth factor A , also known as VEGF-A or Vascular permeability factor. VEGF-A is a 206 amino acid ~45 kDa disulphide linked homodimeric heparin-binding cytokine and member of the PDGF/VEGF growth factor family, which exists as several splice variants of the VEGF gene, and is a potent stimulator of angiogenesis of both normal and cancerous cells, and acts as a regulator of vasculogenesis. The functional forms of VEGF are classified according to the number of amino acid residues within a constitutive polypeptide chain e.g VEGF ₁₆₅ , and all VEGF family members act through binding to the tyrosine kinase receptors VEGFR-1 (Flt-1) and VEGFR-2 (KDR/CD309), whilst the closely-related VEGF-C and VEGF-D, can also bind to		
	recognizes human Vascular endothelial growth factor A , also known as VEGF-A or Vascular permeability factor. VEGF-A is a 206 amino acid ~45 kDa disulphide linked homodimeric heparin-binding cytokine and member of the PDGF/VEGF growth factor family, which exists as several splice variants of the VEGF gene, and is a potent stimulator of angiogenesis of both normal and cancerous cells, and acts as a regulator of vasculogenesis. The functional forms of VEGF are classified according to the number of amino acid residues within a constitutive polypeptide chain e.g VEGF ₁₆₅ , and all VEGF family members act through binding to the tyrosine kinase receptors VEGFR-1 (FIt-1) and VEGFR-2 (KDR/CD309), whilst the closely-related VEGF-C and VEGF-D, can also bind to VEGFR-3 (FIt-4).		
ELISA	recognizes human Vascular endothelial growth factor A , also known as VEGF-A or Vascular permeability factor. VEGF-A is a 206 amino acid ~45 kDa disulphide linked homodimeric heparin-binding cytokine and member of the PDGF/VEGF growth factor family, which exists as several splice variants of the VEGF gene, and is a potent stimulator of angiogenesis of both normal and cancerous cells, and acts as a regulator of vasculogenesis. The functional forms of VEGF are classified according to the number of amino acid residues within a constitutive polypeptide chain e.g VEGF ₁₆₅ , and all VEGF family members act through binding to the tyrosine kinase receptors VEGFR-1 (FIt-1) and VEGFR-2 (KDR/CD309), whilst the closely-related VEGF-C and VEGF-D, can also bind to VEGFR-3 (FIt-4). MCA5750Z may be used in a sandwich ELISA with <u>AHP831B</u> as the detection antibody and <u>PHP091</u> as the standard. MCA5750Z detects a band of approximately 38.2kDa using recombinant human VEGF ₁₆₅		

	recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.			
Guarantee	12 months from date of despatch			
Health And Safety Information	Material Safety Datasheet documentation #10294 available at: https://www.bio-rad-antibodies.com/SDS/MCA5750Z 10294			
Regulatory	For research purposes only			

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77)	HRP
Rabbit Anti Mouse IgG (STAR12)	RPE
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>
Goat Anti Mouse IgG IgA IgM (STAR87) <u>Alk. Phos.</u> , <u>HRP</u>
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>
Goat Anti Mouse IgG (STAR76)	RPE
Rabbit Anti Mouse IgG (STAR13)	HRP
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,
	DyLight®650, DyLight®680, DyLight®800,
	<u>FITC, HRP</u>
Goat Anti Mouse IgG (Fc) (STAR120)	<u>FITC, HRP</u>

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rac	l.com	Email: antibody_sales_uk@bio-rac	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M402158:220718'

Printed on 29 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint