

Datasheet: MCA5775GA BATCH NUMBER 148431

Description:	MOUSE ANTI HUMAN ACTIN BETA
Specificity:	ACTIN BETA
Other names:	ACTB
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	4C2
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>					
	rad-antibodies.com/proto		N			
	luono un obietale m.c. Engenera	Yes	No	Not Determined	Suggested Dilution	
	Immunohistology - Frozen	-			1/50 - 1/100	
	Immunohistology - Paraffin					
	ELISA			•		
	Immunoprecipitation	-		•		
	Western Blotting	-				
	Immunofluorescence Where this antibody has not been tested for use in a particular technique this does not 					
	a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.					
Target Species	Human					
Target Species Species Cross Reactivity	Human Reacts with: Rabbit, Pig, N.B. Antibody reactivity a reactivity is derived from personal communications further information.	and worki testing w	ing conditi /ithin our la	aboratories, peer-revie	wed publications or	
Species Cross	Reacts with: Rabbit, Pig, N.B. Antibody reactivity a reactivity is derived from personal communication	and worki testing w	ing conditi /ithin our la	aboratories, peer-revie	wed publications or	

	supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Synthetic peptide corresponding to the acetylated N-terminal nonapeptide of beta-cytoplasmic actin, coupled to KLH.
External Database Links	UniProt: P60709 Related reagents Entrez Gene: 60 ACTB Related reagents
RRID	AB_2571580
Specificity	 Mouse anti Human actin beta, clone 4C2, recognizes cytoplasmic actin beta (known also as ACTB and Beta Actin), a 41.7 kDa cytoskeletal protein. Actins are highly conserved proteins that are involved in cell motility, structure and integrity and are ubiquitously expressed in all eukaryotic cells. Six different isoforms of actin have been identified in mammals (Vandekerckhove <i>et al.</i> 1978). Clone 4C2 is highly specific for cytoplasmic actin β and does not react with other actin isoforms. Mouse anti Human actin beta, clone 4C2 has been used to look at the remodeling of the actin cytoskeleton following sendai virus infection and the specific role of the viral M protein in actin remodeling (Miazza <i>et al.</i> 2011).
References	 Dugina, V. <i>et al.</i> (2009) Beta and gamma-cytoplasmic actins display distinct distribution and functional diversity. J Cell Sci. 122 (Pt 16): 2980-8. Brockmann, C. <i>et al.</i> (2011) Beta- and gamma-cytoplasmic actins are required for meiosis in mouse oocytes. Biol Reprod. 85 (5): 1025-39. Miazza, V. <i>et al.</i> (2011) Sendai virus induced cytoplasmic actin remodeling correlates with efficient virus particle production. Virology. 410 (1): 7-16. Baranwal, S. <i>et al.</i> (2012) Nonredundant roles of cytoplasmic β- and γ-actin isoforms in regulation of epithelial apical junctions. Mol Biol Cell. 23 (18): 3542-53. Shagieva, G.S. <i>et al.</i> (2012) Actin isoforms and reorganization of adhesion junctions in epithelial-to-mesenchymal transition of cervical carcinoma cells. Biochemistry (Mosc). 77 (11): 1266-76. Dugina, V. <i>et al.</i> (2015) Tumor promotion by γ and suppression by β non-muscle actin isoforms. Oncotarget. 6 (16): 14556-71.

	7. Rubtsova, S.N. et al. (2015) A Novel Role of E-Cadherin-Based Adherens Junctions in			
	Neoplastic Cell Dissemination. PLoS One. 10 (7): e0133578.			
	8. Simiczyjew A et al. (2015) Active invadopodia of mesenchymally migrating cancer cells			
	contain both β and γ cytoplasmic actin isoforms. Exp Cell Res. pii:			
	<u>S0014-4827(15)30146-4.</u>			
	9. Simiczyjew, A. et al. (2017) Involvement of β - and γ -actin isoforms in actin cytoskeleton			
	organization and migration abilities of bleb-forming human colon cancer cells. <u>PLoS One.</u>			
	<u>12 (3): e0173709.</u>			
	10. Novikova, M.V. et al. (2017) [A change in the expression of membrane-associated			
	proteins and cytoplasmic actin isoforms in the progression of human colon tumors]. <u>Arkh</u>			
	Patol. 79 (2): 15-21.			
	11. Chaponnier, C. & Gabbiani, G. (2016) Monoclonal antibodies against muscle actin			
	isoforms: epitope identification and analysis of isoform expression by immunoblot and			
	immunostaining in normal and regenerating skeletal muscle. <u>F1000Res. 5: 416.</u>			
	12. Chen, A. <i>et al.</i> (2017) Cytokinesis requires localized β -actin filament production by an			
	actin isoform specific nucleator. <u>Nat Commun. 8 (1): 1530.</u>			
Further Reading	1. Perrin, B.J. & Ervasti, J.M. (2010) The actin gene family: function follows isoform.			
Further Reading	1. Perrin, B.J. & Ervasti, J.M. (2010) The actin gene family: function follows isoform. <u>Cytoskeleton (Hoboken). 67 (10): 630-4.</u>			
Further Reading Storage	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred.			
	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended.			
	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may			
	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend			
	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may			
	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend			
Storage Guarantee	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch			
Storage Guarantee Health And Safety	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch Material Safety Datasheet documentation #10040 available at:			
Storage Guarantee	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA5775GA			
Storage Guarantee Health And Safety	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch Material Safety Datasheet documentation #10040 available at:			
Storage Guarantee Health And Safety	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA5775GA			

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>
Goat Anti Mouse IgG (STAR76)	RPE
Rabbit Anti Mouse IgG (STAR13)	HRP
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,
	DyLight®650, DyLight®680, DyLight®800,
	FITC, HRP
Rabbit Anti Mouse IgG (STAR9)	FITC
Goat Anti Mouse IgG (STAR77)	HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody sales us@bio-ra	d.com	Email: antibody sales uk@bio-ra	d.com	Email: antibody sales de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368378:200529'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint