

# Datasheet: MCA5775GA BATCH NUMBER 154012

Description:	MOUSE ANTI HUMAN ACTIN BETA
Specificity:	ACTIN BETA
Other names:	ACTB
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	4C2
Isotype:	lgG1
Quantity:	0.1 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is						
	derived from testing withi	derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.					
	communications from the						
	information. For general						
	rad-antibodies.com/proto						
		Yes	No	Not Determined	Suggested Dilution		
	Immunohistology - Frozen	-			1/50 - 1/100		
	Immunohistology - Paraffin			-			
	ELISA						
	Immunoprecipitation						
	Western Blotting	-					
	Immunofluorescence						
	necessarily exclude its us	se in suc	h procedu		ng dilutions are given as		
	•	se in suc nended th	h procedu nat the use	res. Suggested workin er titrates the antibody	ng dilutions are given as		
Target Species	necessarily exclude its us a guide only. It is recomn	se in suc nended th	h procedu nat the use	res. Suggested workin er titrates the antibody	ng dilutions are given as		
	necessarily exclude its us a guide only. It is recomn system using appropriate	se in suc nended th negative	h procedu nat the use e/positive o	res. Suggested workin er titrates the antibody	ng dilutions are given a		
Species Cross	necessarily exclude its us a guide only. It is recomn system using appropriate Human	se in suc nended th negative Mouse, I	h procedu nat the use e/positive o Rat	res. Suggested workin er titrates the antibody controls.	ng dilutions are given as		
Species Cross	necessarily exclude its us a guide only. It is recommon system using appropriate Human Reacts with: Rabbit, Pig,	se in suc nended th negative Mouse, I and worki	h procedu nat the use e/positive e/positive Rat ng conditi	res. Suggested workin er titrates the antibody controls.	ng dilutions are given a for use in their own		
Species Cross	necessarily exclude its us a guide only. It is recommon system using appropriate Human Reacts with: Rabbit, Pig, <b>N.B.</b> Antibody reactivity a reactivity is derived from	se in suc nended th negative Mouse, I and worki testing w	h procedu nat the use e/positive o Rat ng conditi rithin our la	res. Suggested workin er titrates the antibody controls. ons may vary betweer aboratories, peer-revie	ng dilutions are given a for use in their own n species. Cross wed publications or		
Target Species Species Cross Reactivity	necessarily exclude its us a guide only. It is recomm system using appropriate Human Reacts with: Rabbit, Pig, <b>N.B.</b> Antibody reactivity a	se in suc nended th negative Mouse, I and worki testing w	h procedu nat the use e/positive o Rat ng conditi rithin our la	res. Suggested workin er titrates the antibody controls. ons may vary betweer aboratories, peer-revie	ng dilutions are given as for use in their own n species. Cross wed publications or		
Species Cross	necessarily exclude its us a guide only. It is recomm system using appropriate Human Reacts with: Rabbit, Pig, <b>N.B.</b> Antibody reactivity a reactivity is derived from personal communications	se in suc nended th negative Mouse, I and worki testing w	h procedu nat the use e/positive o Rat ng conditi rithin our la	res. Suggested workin er titrates the antibody controls. ons may vary betweer aboratories, peer-revie	ng dilutions are given as for use in their own n species. Cross wed publications or		

	supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Synthetic peptide corresponding to the acetylated N-terminal nonapeptide of beta-cytoplasmic actin, coupled to KLH.
External Database Links	UniProt:         P60709       Related reagents         Entrez Gene:         60       ACTB       Related reagents
RRID	AB_2571580
Specificity	<ul> <li>Mouse anti Human actin beta, clone 4C2, recognizes cytoplasmic actin beta (known also as ACTB and Beta Actin), a 41.7 kDa cytoskeletal protein.</li> <li>Actins are highly conserved proteins that are involved in cell motility, structure and integrity and are ubiquitously expressed in all eukaryotic cells. Six different isoforms of actin have been identified in mammals (Vandekerckhove <i>et al.</i> 1978).</li> <li>Clone 4C2 is highly specific for cytoplasmic actin β and does not react with other actin isoforms. Mouse anti Human actin beta, clone 4C2 has been used to look at the remodeling of the actin cytoskeleton following sendai virus infection and the specific role of the viral M protein in actin remodeling (Miazza <i>et al.</i> 2011).</li> </ul>
References	<ol> <li>Dugina, V. <i>et al.</i> (2009) Beta and gamma-cytoplasmic actins display distinct distribution and functional diversity. J Cell Sci. 122 (Pt 16): 2980-8.</li> <li>Brockmann, C. <i>et al.</i> (2011) Beta- and gamma-cytoplasmic actins are required for meiosis in mouse oocytes. Biol Reprod. 85 (5): 1025-39.</li> <li>Miazza, V. <i>et al.</i> (2011) Sendai virus induced cytoplasmic actin remodeling correlates with efficient virus particle production. Virology. 410 (1): 7-16.</li> <li>Baranwal, S. <i>et al.</i> (2012) Nonredundant roles of cytoplasmic β- and γ-actin isoforms in regulation of epithelial apical junctions. Mol Biol Cell. 23 (18): 3542-53.</li> <li>Shagieva, G.S. <i>et al.</i> (2012) Actin isoforms and reorganization of adhesion junctions in epithelial-to-mesenchymal transition of cervical carcinoma cells. Biochemistry (Mosc). 77 (11): 1266-76.</li> <li>Dugina, V. <i>et al.</i> (2015) Tumor promotion by γ and suppression by β non-muscle actin isoforms. Oncotarget. 6 (16): 14556-71.</li> </ol>

	7. Rubtsova, S.N. et al. (2015) A Novel Role of E-Cadherin-Based Adherens Junctions in
	Neoplastic Cell Dissemination. PLoS One. 10 (7): e0133578.
	8. Simiczyjew A et al. (2015) Active invadopodia of mesenchymally migrating cancer cells
	contain both $\beta$ and $\gamma$ cytoplasmic actin isoforms. Exp Cell Res. pii:
	<u>S0014-4827(15)30146-4.</u>
	9. Simiczyjew, A. et al. (2017) Involvement of $\beta$ - and $\gamma$ -actin isoforms in actin cytoskeleton
	organization and migration abilities of bleb-forming human colon cancer cells. <u>PLoS One.</u>
	<u>12 (3): e0173709.</u>
	10. Novikova, M.V. et al. (2017) [A change in the expression of membrane-associated
	proteins and cytoplasmic actin isoforms in the progression of human colon tumors]. <u>Arkh</u>
	Patol. 79 (2): 15-21.
	11. Chaponnier, C. & Gabbiani, G. (2016) Monoclonal antibodies against muscle actin
	isoforms: epitope identification and analysis of isoform expression by immunoblot and
	immunostaining in normal and regenerating skeletal muscle. <u>F1000Res. 5: 416.</u>
	12. Chen, A. <i>et al.</i> (2017) Cytokinesis requires localized $\beta$ -actin filament production by an
	actin isoform specific nucleator. <u>Nat Commun. 8 (1): 1530.</u>
Further Reading	1. Perrin, B.J. & Ervasti, J.M. (2010) The actin gene family: function follows isoform.
Further Reading	1. Perrin, B.J. & Ervasti, J.M. (2010) The actin gene family: function follows isoform. <u>Cytoskeleton (Hoboken). 67 (10): 630-4.</u>
Further Reading Storage	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred.
	Cytoskeleton (Hoboken). 67 (10): 630-4.         Store at +4°C or at -20°C if preferred.         Storage in frost-free freezers is not recommended.
	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may
	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend
	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may
	Cytoskeleton (Hoboken). 67 (10): 630-4. Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend
Storage Guarantee	Cytoskeleton (Hoboken). 67 (10): 630-4.         Store at +4°C or at -20°C if preferred.         Storage in frost-free freezers is not recommended.         This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.         12 months from date of despatch
Storage Guarantee Health And Safety	Cytoskeleton (Hoboken). 67 (10): 630-4.         Store at +4°C or at -20°C if preferred.         Storage in frost-free freezers is not recommended.         This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.         12 months from date of despatch         Material Safety Datasheet documentation #10040 available at:
Storage Guarantee	Cytoskeleton (Hoboken). 67 (10): 630-4.         Store at +4°C or at -20°C if preferred.         Storage in frost-free freezers is not recommended.         This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.         12 months from date of despatch         Material Safety Datasheet documentation #10040 available at:         https://www.bio-rad-antibodies.com/SDS/MCA5775GA
Storage Guarantee Health And Safety	Cytoskeleton (Hoboken). 67 (10): 630-4.         Store at +4°C or at -20°C if preferred.         Storage in frost-free freezers is not recommended.         This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.         12 months from date of despatch         Material Safety Datasheet documentation #10040 available at:
Storage Guarantee Health And Safety	Cytoskeleton (Hoboken). 67 (10): 630-4.         Store at +4°C or at -20°C if preferred.         Storage in frost-free freezers is not recommended.         This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.         12 months from date of despatch         Material Safety Datasheet documentation #10040 available at:         https://www.bio-rad-antibodies.com/SDS/MCA5775GA

# **Related Products**

### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87	) <u>HRP</u>
Goat Anti Mouse IgG (STAR76)	RPE
Rabbit Anti Mouse IgG (STAR13)	HRP
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,
	DyLight®650, DyLight®680, DyLight®800,
	FITC, HRP
Rabbit Anti Mouse IgG (STAR9)	FITC
Goat Anti Mouse IgG (STAR77)	HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody sales us@bio-ra	d.com	Email: antibody sales uk@bio-ra	d.com	Email: antibody sales de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368378:200529'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint