

Datasheet: MCA5775GA BATCH NUMBER 164278

Description:	MOUSE ANTI HUMAN ACTIN BETA
Specificity:	ACTIN BETA
Other names:	ACTB
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	4C2
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications		This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further					
	e e e						
		-					
	•	information. For general protocol recommendations, please visit <u>www.bio-</u>					
	rad-antibodies.com/proto	rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution		
	Immunohistology - Frozen	-			1/50 - 1/100		
	Immunohistology - Paraffin						
	ELISA			-			
	Immunoprecipitation			•			
	Western Blotting	-					
	Immunofluorescence	-					
	necessarily exclude its us a guide only. It is recomn	se in suc	h procedu		ng dilutions are given as		
	necessarily exclude its us	se in suc nended th	h procedu nat the use	res. Suggested workin er titrates the antibody	ng dilutions are given as		
Target Species	necessarily exclude its us a guide only. It is recomn	se in suc nended th	h procedu nat the use	res. Suggested workin er titrates the antibody	ng dilutions are given as		
Target Species Species Cross	necessarily exclude its us a guide only. It is recomn system using appropriate	se in suc nended th negative	h procedu nat the use e/positive	res. Suggested workin er titrates the antibody	ng dilutions are given as		
	necessarily exclude its us a guide only. It is recomn system using appropriate Human	se in suc nended th negative Mouse, I	h procedu nat the use e/positive Rat	res. Suggested workin er titrates the antibody controls.	ng dilutions are given as r for use in their own		
Species Cross	necessarily exclude its us a guide only. It is recomm system using appropriate Human Reacts with: Rabbit, Pig, N.B. Antibody reactivity a	se in suc nended th negative Mouse, I and worki	h procedu nat the use e/positive Rat ng conditi	res. Suggested workin er titrates the antibody controls.	ng dilutions are given as for use in their own n species. Cross		
Species Cross	necessarily exclude its us a guide only. It is recommon system using appropriate Human Reacts with: Rabbit, Pig, N.B. Antibody reactivity a reactivity is derived from	se in suc nended th negative Mouse, I and worki testing w	h procedu nat the use e/positive Rat ng conditi rithin our la	res. Suggested workin er titrates the antibody controls. ons may vary betweer aboratories, peer-revie	ng dilutions are given as for use in their own n species. Cross ewed publications or		
Species Cross	necessarily exclude its us a guide only. It is recomm system using appropriate Human Reacts with: Rabbit, Pig, N.B. Antibody reactivity a	se in suc nended th negative Mouse, I and worki testing w	h procedu nat the use e/positive Rat ng conditi rithin our la	res. Suggested workin er titrates the antibody controls. ons may vary betweer aboratories, peer-revie	ng dilutions are given as for use in their own n species. Cross ewed publications or		
Species Cross	necessarily exclude its us a guide only. It is recomm system using appropriate Human Reacts with: Rabbit, Pig, N.B. Antibody reactivity a reactivity is derived from personal communications	se in suc nended th negative Mouse, I and worki testing w	h procedu nat the use e/positive Rat ng conditi rithin our la	res. Suggested workin er titrates the antibody controls. ons may vary betweer aboratories, peer-revie	ng dilutions are given as for use in their own n species. Cross ewed publications or		

	supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Synthetic peptide corresponding to the acetylated N-terminal nonapeptide of beta-cytoplasmic actin, coupled to KLH.
External Database Links	UniProt: P60709 Related reagents Entrez Gene: 60 ACTB Related reagents
RRID	AB_2571580
Specificity	 Mouse anti Human actin beta, clone 4C2, recognizes cytoplasmic actin beta (known also as ACTB and Beta Actin), a 41.7 kDa cytoskeletal protein. Actins are highly conserved proteins that are involved in cell motility, structure and integrity and are ubiquitously expressed in all eukaryotic cells. Six different isoforms of actin have been identified in mammals (Vandekerckhove <i>et al.</i> 1978).
	Clone 4C2 is highly specific for cytoplasmic actin β and does not react with other actin isoforms. Mouse anti Human actin beta, clone 4C2 has been used to look at the remodeling of the actin cytoskeleton following sendai virus infection and the specific role of the viral M protein in actin remodeling (<u>Miazza <i>et al.</i> 2011</u>).
References	 Brockmann, C. <i>et al.</i> (2011) Beta- and gamma-cytoplasmic actins are required for meiosis in mouse oocytes. <u>Biol Reprod. 85 (5): 1025-39.</u> Miazza, V. <i>et al.</i> (2011) Sendai virus induced cytoplasmic actin remodeling correlates with efficient virus particle production. <u>Virology. 410 (1): 7-16.</u> Baranwal, S. <i>et al.</i> (2012) Nonredundant roles of cytoplasmic β- and γ-actin isoforms in regulation of epithelial apical junctions. <u>Mol Biol Cell. 23 (18): 3542-53.</u> Shagieva, G.S. <i>et al.</i> (2012) Actin isoforms and reorganization of adhesion junctions in epithelial-to-mesenchymal transition of cervical carcinoma cells. <u>Biochemistry (Mosc). 77 (11): 1266-76.</u> Dugina, V. <i>et al.</i> (2015) Tumor promotion by γ and suppression by β non-muscle actin isoforms. <u>Oncotarget. 6 (16): 14556-71.</u> Rubtsova, S.N. <i>et al.</i> (2015) A Novel Role of E-Cadherin-Based Adherens Junctions in Neoplastic Cell Dissemination. <u>PLoS One. 10 (7): e0133578.</u>

	7. Simiczyjew, A. <i>et al.</i> (2015) Active invadopodia of mesenchymally migrating cancer
	cells contain both β and γ cytoplasmic actin isoforms. Exp Cell Res. 339 (2): 206-19.
	8. Simiczyjew, A. <i>et al.</i> (2017) Involvement of β - and γ -actin isoforms in actin cytoskeleton organization and migration abilities of bleb-forming human colon cancer cells. <u>PLoS One.</u> <u>12 (3): e0173709.</u>
	9. Novikova, M.V. et al. (2017) [A change in the expression of membrane-associated
	proteins and cytoplasmic actin isoforms in the progression of human colon tumors]. <u>Arkh</u> Patol. 79 (2): 15-21.
	10. Chaponnier, C. & Gabbiani, G. (2016) Monoclonal antibodies against muscle actin isoforms: epitope identification and analysis of isoform expression by immunoblot and immunostaining in normal and regenerating skeletal muscle. F1000Res. 5: 416.
	11. Chen, A. <i>et al.</i> (2017) Cytokinesis requires localized β-actin filament production by an actin isoform specific nucleator. <u>Nat Commun. 8 (1): 1530.</u>
	12. Dugina, V. <i>et al.</i> (2021) Impaired Expression of Cytoplasmic Actins Leads to Chromosomal Instability of MDA-MB-231 Basal-Like Mammary Gland Cancer Cell Line. <u>Molecules. 26(8):2151.</u>
	Molecules. 20(0).2131.
Further Reading	1. Perrin, B.J. & Ervasti, J.M. (2010) The actin gene family: function follows isoform. Cytoskeleton (Hoboken). 67 (10): 630-4.
Further Reading Storage	
	Cytoskeleton (Hoboken). 67 (10): 630-4. This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in
	Cytoskeleton (Hoboken). 67 (10): 630-4. This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Cytoskeleton (Hoboken). 67 (10): 630-4. This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in
Storage	Cytoskeleton (Hoboken). 67 (10): 630-4. This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77)	<u>HRP</u>
Rabbit Anti Mouse IgG (STAR12)	<u>RPE</u>
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>
Goat Anti Mouse IgG IgA IgM (STAR87) <u>Alk. Phos.</u> , <u>HRP</u>
Goat Anti Mouse IgG (STAR76)	<u>RPE</u>
Goat Anti Mouse IgG (Fc) (STAR120)	<u>FITC, HRP</u>
Rabbit Anti Mouse IgG (STAR13)	<u>HRP</u>
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

Alk. Phos., DyLight®488, DyLight®550, DyLight®650, DyLight®680, DyLight®800, FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide
America	Fax: +1 919 878 3751	
	Email: antibody_sales_us@bio-rad	.com

 Tel: +44 (0)1865 852 700
 Europe

 Fax: +44 (0)1865 852 739
 Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M381760:210512'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint