

# Datasheet: MCA5782F BATCH NUMBER 0112

Description:	MOUSE ANTI HUMAN SIGLEC-7:FITC
Specificity:	SIGLEC-7
Other names:	CD328
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	S7.7
Isotype:	lgG1
Quantity:	0.1 mg

# **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	-			Neat - 1/10		
	Functional Assays (1)			•			
	Where this antibody has not been tested for use in a particular technique this does not						
	necessarily exclude i	ts use in such p	rocedures	. Suggested workin	g dilutions are given as		
	a guide only. It is reco	a guide only. It is recommended that the user titrates the antibody for use in their own					
	system using approp	riate negative/po	ositive con	trols.			
	(1) Bio-Rad recommends the use of <u>MCA5782EL</u> for functional assays.						
Target Species	Human						
Species Cross	Reacts with: Rhesus	Monkey					
Reactivity	<b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	Fluorophore	Excitation Ma	x (nm) Er	mission Max (nm)			
	FITC	490		525			
Preparation	Purified IgG prepared	l by affinity chro	matograpl	hy on Protein G fror	n tissue culture		

	supernatant			
Buffer Solution	Phosphate buffered saline.			
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin			
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml			
Immunogen	Siglec-7 3T3 transfected cells.			
External Database Links	UniProt: Q9Y286 Related reagents Entrez Gene:			
	27036 SIGLEC7 Related reagents			
Synonyms	AIRM1			
Fusion Partners	Spleen cells from immunised Balb/c mice, were fused with cells of the Sp2/0 myeloma cell line.			
Specificity	Mouse anti Human Siglec-7 antibody, cloneS7.7 recognizes human Sialic acid-binding Ig-like lectin 7, otherwise known as CD328, a putative adhesion molecule and member of the Ig superfamily, predominantly expressed by both resting and activated natural killer (NK) cells, and also at lower levels by monocytes, granulocytes, monocyte-derived macrophages and dendritic cells, and a small population of CD8+ memory T cells. Classed as a CD33-related Siglec, Siglec-7 preferentially binds to glycoconjugates containing alpha-2,3- or alpha-2,6-linked sialic acid, and acts as a novel inhibitory receptor			
	for NK cells, mediating the inhibition of NK cell cytotoxicity, and implicated in the regulation of NK cell and T cell activation, and hematopoiesis.			
	Mouse anti Human Siglec-7 antibody, cloneS7.7 is a blocking antibody ( <u>Avril <i>et al.</i> 2006</u> ), and has been shown to recognize Siglec-7 in Rhesus monkey ( <u>Jaroenpool <i>et al.</i> 2007</u> ).			
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells.			
References	<ol> <li>Nicoll, G. <i>et al.</i> (1999) Identification and characterization of a novel siglec, siglec-7, expressed by human natural killer cells and monocytes. J Biol Chem. 274 (48): 34089-95.</li> <li>Jaroenpool, J. <i>et al.</i> (2007) Differences in the constitutive and SIV infection induced expression of Siglecs by hematopoietic cells from non-human primates. <u>Cell Immunol. 250</u> (1-2): 91-104.</li> <li>Avril T <i>et al.</i> (2006) Sialic acid-binding immunoglobulin-like lectin 7 mediates selective recognition of sialylated glycans expressed on <i>Campylobacter jejuni</i> lipooligosaccharides. <u>Infect Immun. 74 (7): 4133-41.</u></li> <li>Nguyen, D.H. <i>et al.</i> (2006) Loss of Siglec expression on T lymphocytes during human</li> </ol>			

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA5782F 10041
Regulatory	For research purposes only

## **Related Products**

### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL: FITC (MCA928F)

### **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852
	Email: antibody_sales_	us@bio-rad.com	Email: antibody_sales

H: +44 (0)1865 852 700 **Europe** ax: +44 (0)1865 852 739 mail: antibody\_sales\_uk@bio-rad.com Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368390:200529'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint