

Datasheet: MCA5782F

BATCH NUMBER 162821

Description:	MOUSE ANTI HUMAN SIGLEC-7:FITC
Specificity:	SIGLEC-7
Other names:	CD328
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	S7.7
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Functional Assays (1)			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **Bio-Rad recommends the use of [MCA5782EL](#) for functional assays.**

Target Species

Human

Species Cross Reactivity

Reacts with: Rhesus Monkey

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture

supernatant

Buffer Solution Phosphate buffered saline.

Preservative 0.09% Sodium Azide (NaN₃)
Stabilisers 1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.1 mg/ml

Immunogen Siglec-7 3T3 transfected cells.

External Database Links

UniProt:

[Q9Y286](#)

[Related reagents](#)

Entrez Gene:

[27036](#)

SIGLEC7

[Related reagents](#)

Synonyms AIRM1

Fusion Partners Spleen cells from immunized Balb/c mice, were fused with cells of the Sp2/0 myeloma cell line.

Specificity **Mouse anti Human Siglec-7 antibody, cloneS7.7** recognizes human Sialic acid-binding Ig-like lectin 7, otherwise known as CD328, a putative adhesion molecule and member of the Ig superfamily, predominantly expressed by both resting and activated natural killer (NK) cells, and also at lower levels by monocytes, granulocytes, monocyte-derived macrophages and dendritic cells, and a small population of CD8+ memory T cells.

Classed as a CD33-related Siglec, Siglec-7 preferentially binds to glycoconjugates containing alpha-2,3- or alpha-2,6-linked sialic acid, and acts as a novel inhibitory receptor for NK cells, mediating the inhibition of NK cell cytotoxicity, and implicated in the regulation of NK cell and T cell activation, and hematopoiesis.

Mouse anti Human Siglec-7 antibody, cloneS7.7 is a blocking antibody ([Avril *et al.* 2006](#)), and has been shown to recognize Siglec-7 in Rhesus monkey ([Jaroenpool *et al.* 2007](#)).

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells.

References

1. Nicoll, G. *et al.* (1999) Identification and characterization of a novel siglec, siglec-7, expressed by human natural killer cells and monocytes. [J Biol Chem. 274 \(48\): 34089-95.](#)
2. Nguyen, D.H. *et al.* (2006) Loss of Siglec expression on T lymphocytes during human evolution. [Proc Natl Acad Sci U S A. 103 \(20\): 7765-70.](#)
3. Avril T *et al.* (2006) Sialic acid-binding immunoglobulin-like lectin 7 mediates selective recognition of sialylated glycans expressed on *Campylobacter jejuni* lipooligosaccharides. [Infect Immun. 74 \(7\): 4133-41.](#)
4. Jaroenpool, J. *et al.* (2007) Differences in the constitutive and SIV infection induced expression of Siglecs by hematopoietic cells from non-human primates. [Cell Immunol. 250](#)

[\(1-2\): 91-104.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA5782F>
10041

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M384606:210513'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)