

Datasheet: MCA5783GA

Description:	MOUSE ANTI HUMAN SIGLEC-10
Specificity:	SIGLEC-10
Other names:	SIALIC ACID-BINDING IG-LIKE LECTIN 10
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	5G6
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/25 - 1/200
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant human Siglec-10, fused with the Fc region of human IgG

**External Database
Links**

UniProt:

[Q96LC7](#) [Related reagents](#)

Entrez Gene:

[89790](#) SIGLEC10 [Related reagents](#)

Synonyms

SLG2

Specificity

Mouse anti Human Siglec-10 antibody, clone 5G6 recognises human Siglec-10 (Sialic acid-binding Ig-like lectin 10), a putative adhesion molecule and member of the Ig superfamily, expressed by monocytes, B cells, eosinophils, and at a higher level by a subpopulation of CD16+CD56- natural killer (NK) cells. Structurally, Siglec-10 is most similar to the CD33-related group of Siglecs, and preferentially binds to glycoconjugates containing alpha-2,3- or alpha-2,6-linked sialic acid. Siglec-10 acts as a substrate for VAP-1 (Vascular adhesion protein-1), a glycoprotein expressed on endothelium during inflammation, which is involved in primary amine oxidation and leucocyte trafficking, ([Kivi et al. 2009](#)). This interaction between Siglec-10 and VAP-1, implicates Siglec-10 in endothelial lymphocyte adhesion and in the modulation of the inflammatory microenvironment. Mouse anti Human Siglec-10 antibody, clone 5G6 does not cross-react with Siglecs 3, 5, 7, 8 and 9 ([Munday et al. 2001](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells.

References

1. Munday, J. *et al.* (2001) Identification, characterization and leucocyte expression of Siglec-10, a novel human sialic acid-binding receptor. [Biochem J. 355 \(Pt 2\): 489-97.](#)
2. Kivi, E. *et al.* (2009) Human Siglec-10 can bind to vascular adhesion protein-1 and serves as its substrate. [Blood. 114 \(26\): 5385-92.](#)
3. Nguyen, D.H. *et al.* (2006) Loss of Siglec expression on T lymphocytes during human evolution. [Proc Natl Acad Sci U S A. 103 \(20\): 7765-70.](#)

Further Reading

1. Crocker, P.R. (2005) Siglecs in innate immunity. [Curr Opin Pharmacol. 5 \(4\): 431-7.](#)

Storage

Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

18 months from date of despatch.

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@680](#),
[DyLight@800](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M343935:190111'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)