

Datasheet: MCA5787F BATCH NUMBER 166101

Description:	MOUSE ANTI HUMAN SIGLEC-5/SIGLEC-14:FITC
Specificity:	SIGLEC-5/SIGLEC-14
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	1A5
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat - 1/10
Immunofluorescence				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human			
Species Cross	Reacts with: Chir	mpanzee		
Reactivity	reactivity is derive	ed from testing within our land	ons may vary between species aboratories, peer-reviewed pub ers. Please refer to references i	lications or
Product Form	Purified IgG conju	ugated to Fluorescein Isoth	niocyanate Isomer 1 (FITC) - lic	quid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	FITC	490	525	
Preparation	Purified IgG prep supernatant	ared by affinity chromatog	raphy on Protein A from tissue	culture

Buffer Solution	Phosphate buffered saline	
Preservative	0.09% Sodium Azide (NaN ₃)	
Stabilisers	1% Bovine Serum Albumin	
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml	
Immunogen	Siglec-5-Fc protein, consisting of the full-length extracellular region of human Siglec-5, fused with the Fc region of human IgG1.	
External Database		
Links	UniProt:	
	O15389 Related reagents	
	Q08ET2 Related reagents	
	Entrez Gene:	
	8778 SIGLEC5 Related reagents	
	100049587 SIGLEC14 Related reagents	
Synonyms	CD33L2, OBBP2	
Specificity	Mouse anti Human Siglec-5/Siglec-14 antibody, clone 1A5 recognizes human Siglec-5 (Sialic acid-binding Ig-like lectin 5), otherwise known as CD170, a novel sialic-acid-binding Ig-like lectin, and member of the Ig superfamily, expressed by dendritic cells (DCs), activated macrophages, neutrophils, and cells of the monocyte/myeloid lineage. Mouse anti Human Siglec-5/Siglec-14 antibody, clone 1A5, is one of several Siglec-5	
	antibodies which also recognises human Siglec-14 (<u>Angata et al. 2006</u>). Siglec-14 shares an almost identical sequence with Siglec-5 within the first two Ig-like domains, indicating partial gene conversion between these two Siglecs, also evident in other primate species.	
	Siglec-5 is also related to the myelomonocytic-derived adhesion molecule CD33 (Siglec-3), and mediates sialic-acid dependent binding to cells, as well as acting as an inhibitory receptor in the down-regulation of cell activation.	
	Structurally, Siglec-5 contains an immunoreceptor tyrosine-based inhibitor motif (ITIM), which plays a part in the modulation of cellular responses, and when phosphorylated, can bind to the SH2 domain of several SH2-containing phosphatases. Siglec-14 is a putative sialic-acid binding adhesion molecule, and member of the Ig superfamily, predominantly expressed in hematopoietic tissues, which has been shown to associate with the	

Flow Cytometry

Use 10ul of the suggested working dilution to label 10^6 cells.

1A5 cross reacts with Chimpanzee (Jaroenpool et al. 2007).

References

1. Cornish, A.L. *et al.* (1998) Characterization of siglec-5, a novel glycoprotein expressed on myeloid cells related to CD33. <u>Blood. 92 (6): 2123-32.</u>

activating adapter protein DAP12. Mouse anti Human Siglec-5/Siglec-14 antibody, clone

- 2. Avril, T. et al. (2005) Siglec-5 (CD170) can mediate inhibitory signaling in the absence of immunoreceptor tyrosine-based inhibitory motif phosphorylation. J Biol Chem. 280 (20): 19843-51.
- 3. Angata, T. et al. (2006) Discovery of Siglec-14, a novel sialic acid receptor undergoing concerted evolution with Siglec-5 in primates. FASEB J. 20: 1964-1973.
- 4. Nguyen, D.H. et al. (2006) Loss of Siglec expression on T lymphocytes during human evolution. Proc Natl Acad Sci U S A. 103 (20): 7765-70.
- 5. Jaroenpool, J. et al. (2007) Differences in the constitutive and SIV infection induced expression of Siglecs by hematopoietic cells from non-human primates. Cell Immunol. 250 (1-2): 91-104.

Further Reading

1. Crocker, P.R. (2005) Siglecs in innate immunity. Curr Opin Pharmacol. 5 (4): 431-7.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA5787F 10041		
Regulatory	For research purposes only		

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21

America

Fax: +1 919 878 3751

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419907:230628'

Printed on 18 Jan 2024