

Datasheet: MCA5787PE

BATCH NUMBER 0811R

Description:	MOUSE ANTI HUMAN SIGLEC-5/SIGLEC-14:RPE
Specificity:	SIGLEC-5/SIGLEC-14
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	1A5
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human			
Species Cross	Reacts with: Chimp	anzee		
Reactivity	reactivity is derived	from testing within our I	ons may vary between species. Cro aboratories, peer-reviewed publicati ors. Please refer to references indica	ions (
Product Form	Purified IgG conjug	ated to R. Phycoerythrin	(RPE) - lyophilized	
Reconstitution	Reconstitute with 1	.0ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	RPE 488nm laser	496	578	
Preparation	Purified IgG prepar supernatant	ed by affinity chromatog	raphy on Protein G from tissue cultu	ıre

Buffer Solution	Phosphate bu	uffered saline
Preservative	0.09% Sodiu	m Azide (NaN ₃)
Stabilisers	1% Bovine S	erum Albumin
	5% Sucrose	
Immunogen		rotein, consisting of the full-length extracellular region of human Siglec-5, e Fc region of human IgG1.
External Database		
Links	UniProt:	
	<u>O15389</u>	Related reagents
	Q08ET2	Related reagents
	Entrez Gene	s:

8778 SIGLEC5 Related reagents 100049587 SIGLEC14 Related reagents

Synonyms CD33L2, OBBP2

Specificity

Mouse anti Human Siglec-5/Siglec-14 antibody, clone 1A5 recognizes human Siglec-5 (Sialic acid-binding Ig-like lectin 5), otherwise known as CD170, a novel sialic-acid-binding Ig-like lectin, and member of the Ig superfamily, expressed by dendritic cells (DCs), activated macrophages, neutrophils, and cells of the monocyte/myeloid lineage.

Mouse anti Human Siglec-5/Siglec-14 antibody, clone 1A5, is one of several Siglec-5 antibodies which also recognises human Siglec-14 (Angata et al. 2006). Siglec-14 shares an almost identical sequence with Siglec-5 within the first two Ig-like domains, indicating partial gene conversion between these two Siglecs, also evident in other primate species.

Siglec-5 is also related to the myelomonocytic-derived adhesion molecule CD33 (Siglec-3), and mediates sialic-acid dependent binding to cells, as well as acting as an inhibitory receptor in the down-regulation of cell activation.

Structurally, Siglec-5 contains an immunoreceptor tyrosine-based inhibitor motif (ITIM), which plays a part in the modulation of cellular responses, and when phosphorylated, can bind to the SH2 domain of several SH2-containing phosphatases. Siglec-14 is a putative sialic-acid binding adhesion molecule, and member of the Ig superfamily, predominantly expressed in hematopoietic tissues, which has been shown to associate with the activating adapter protein DAP12. Mouse anti Human Siglec-5/Siglec-14 antibody, clone 1A5 cross reacts with Chimpanzee (Jaroenpool et al. 2007).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells.

References

- 1. Cornish, A.L. et al. (1998) Characterization of siglec-5, a novel glycoprotein expressed on myeloid cells related to CD33. Blood. 92 (6): 2123-32.
- 2. Avril, T. et al. (2005) Siglec-5 (CD170) can mediate inhibitory signaling in the absence of immunoreceptor tyrosine-based inhibitory motif phosphorylation. J Biol Chem. 280 (20):

19843-51.

- 3. Nguyen, D.H. *et al.* (2006) Loss of Siglec expression on T lymphocytes during human evolution. <u>Proc Natl Acad Sci U S A. 103 (20): 7765-70.</u>
- 4. Jaroenpool, J. *et al.* (2007) Differences in the constitutive and SIV infection induced expression of Siglecs by hematopoietic cells from non-human primates. <u>Cell Immunol. 250</u> (1-2): 91-104.
- 5. Angata, T. *et al.* (2006) Discovery of Siglec-14, a novel sialic acid receptor undergoing concerted evolution with Siglec-5 in primates. <u>FASEB J. 20: 1964-1973.</u>

Further Reading	1. Crocker, P.R. (2005) Siglecs in innate immunity. Curr Opin Pharmacol. 5 (4): 431-7
Storage	Prior to reconstitution store at +4°C.
	After reconstitution store at +4°C.
	DO NOT FREEZE. This product should be stored undiluted. This product is photoser
	and should be protected from light. Should this product contain a precipitate we
	recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety	Material Safety Datasheet documentation #20487 available at:
Information	https://www.bio-rad-antibodies.com/SDS/MCA5787PE
	20487
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

Email: antibody_sales_us@bio-rad.com

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375605:210104'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint