

# Datasheet: MCA5788GA

**BATCH NUMBER 154533**

<b>Description:</b>	RAT ANTI MOUSE TIM-1
<b>Specificity:</b>	TIM-1
<b>Other names:</b>	T-CELL IMMUNOGLOBULIN MUCIN 1
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	RMT1-10
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/200
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative</b>	0.09% Sodium Azide (NaN <sub>3</sub> )

## Stabilisers

---

Carrier Free	Yes
--------------	-----

---

Approx. Protein Concentrations	IgG concentration 1 mg/ml
--------------------------------	---------------------------

---

Immunogen	Full-length mouse Tim-1-Ig, containing both the IgV and mucin domains of Tim-1.
-----------	---

---

External Database Links	<p><b>UniProt:</b> <a href="#">Q5QNS5</a>   <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b> <a href="#">171283</a>   Havcr1   <a href="#">Related reagents</a></p>
-------------------------	--

---

Synonyms	Kim1, Tim1, Timd1
----------	-------------------

---

Fusion Partners	Lymph node cells from immunised SD rats were fused with cells of the P3U1 myeloma cell line.
-----------------	--

---

Specificity	<p><b>Rat anti Mouse TIM-1 antibody, clone RMT1-10</b> recognizes mouse Tim-1 (T-cell immunoglobulin mucin 1), a cell surface glycoprotein and member of the immunoglobulin superfamily, which was first identified as the hepatitis A virus cellular receptor (HAVCR), and is highly expressed in the liver (isoform Tim-1a) and the kidneys (isoform Tim-1b).</p> <p>Tim-1 is highly expressed by activated CD4<sup>+</sup> T cells, and acts as a positive/negative co-stimulatory molecule of T cell proliferation, cytokine production and tolerance abrogation, and is the receptor for Tim-4, expressed on APCs. Tim-1 expression is greater on Th2 than Th1 cells, and studies have shown that interaction between Tim-1 on Th2 cells and Tim-4 on dendritic cells (DCs), enhances Th2 cell function, and have implicated Tim-1 as a critical player in the development of atopic disease, and in particular airway hypersensitivity.</p> <p>Polymorphic forms of Tim-1 are associated with increased susceptibility to asthma, eczema and Rheumatoid arthritis.</p> <p>Rat anti Mouse TIM-1 antibody, clone RMT1-10 has been shown to promote Th2 responses, and inhibit antigen-specific T cell proliferation, in contrast to the agonistic function of many Tim-1 antibodies (<a href="#">Xiao et al. 2007</a>).</p> <p>Rat anti Mouse TIM-1 antibody, clone RMT1-10 has been shown to reduce the severity of EAE (experimental autoimmune encephalomyelitis) and delay disease onset, in mice studies (<a href="#">Xiao et al. 2007</a>).</p>
-------------	---

---

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
----------------	---

---

References	1. Xiao, S. et al. (2007) Differential engagement of Tim-1 during activation can positively or negatively costimulate T cell expansion and effector function. <a href="#">J Exp Med. 204 (7):</a>
------------	---

[1691-702.](#)

2. Yuan, X. *et al.* (2009) Targeting Tim-1 to overcome resistance to transplantation tolerance mediated by CD8 T17 cells. [Proc Natl Acad Sci U S A. 106 \(26\): 10734-9.](#)
3. Ueno, T. *et al.* (2008) The emerging role of T cell Ig mucin 1 in alloimmune responses in an experimental mouse transplant model. [J Clin Invest. 118: 742-51.](#)
4. Fukushima, A. *et al.* (2007) Antibodies to T-cell Ig and mucin domain-containing proteins (Tim)-1 and -3 suppress the induction and progression of murine allergic conjunctivitis. [Biochem Biophys Res Commun. 353: 211-6.](#)
5. Rong, S. *et al.* (2011) The TIM-1:TIM-4 pathway enhances renal ischemia-reperfusion injury. [J Am Soc Nephrol. 22: 484-95.](#)
6. Arai, S. *et al.* (2016) Apoptosis inhibitor of macrophage protein enhances intraluminal debris clearance and ameliorates acute kidney injury in mice. [Nat Med. 22 \(2\): 183-93.](#)

---

**Further Reading**

1. Freeman, G.J. *et al.* (2010) TIM genes: a family of cell surface phosphatidylserine receptors that regulate innate and adaptive immunity. [Immunol Rev. 235 \(1\): 172-89.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA5788GA>  
10040

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	<a href="#">DyLight®800</a>
Rabbit Anti Rat IgG (STAR17...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR72...)	<a href="#">HRP</a>
Goat Anti Rat IgG (STAR69...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR73...)	<a href="#">RPE</a>
Rabbit Anti Rat IgG (STAR21...)	<a href="#">HRP</a>
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	<a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®800</a>
Goat Anti Rat IgG (STAR131...)	<a href="#">Alk. Phos.</a> , <a href="#">Biotin</a>

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>	<b>To find a</b>
----------------------------------	---	------------------	---	---------------	---	------------------

batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M368402:200529'

Printed on 18 Jan 2024

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)