

Datasheet: MCA5799GA

Description:	MOUSE ANTI RAT CD1d	
Specificity:	CD1d	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	WTH1	
Isotype:	IgG2a	
Quantity:	0.1 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/2,000
Immunohistology - Frozen	•			
Immunohistology - Paraffin			•	
ELISA	•			
Immunoprecipitation	•			
Western Blotting (1)	•			
Functional Assays (2)	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

- (1) Clone WTH1 recognizes CD1d under non-reducing conditions (Monzon-Casanova, E. et al.).
- (2) This antibody contains sodium azide, removal by dialysis is recommended prior to use in functional assays.

Target Species	Rat
Species Cross	Reacts with: Mouse
Reactivity	Does not react with:Cotton rat
	N.B. Antibody reactivity and working conditions may vary between sp

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Purified IgG - liquid			
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant			
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)			
Approx. Protein Concentrations	IgG concentration 0.5mg/ml			
Immunogen	Rat CD1d-transduced M12.4.1.C3 cells.			
External Database Links	UniProt: P11609 Related reagents Q63493 Related reagents			
	Entrez Gene:			
	<u>12479</u> Cd1d1 <u>Related reagents</u> <u>25109</u> Cd1d1 <u>Related reagents</u>			
Synonyms	Cd1.1, Cd1d1			
RRID	AB_10922750			
Specificity	Mouse anti Rat CD1d antibody, clone WTH1 recognizes rat CD1d, a glycoprotein expressed on cortical thymocytes, certain T-cell leukaemias and various other antigen presenting cells. CD1d binds self and non-self glycolipids and presents them to T-cell receptors on natural killer cells. When activated, natural killer cells produce Th1 and Th2 cytokines. Clone WTH1 binds with high affinity to the mature CD1d heavy chain independently of beta-2-microglobulin. Mouse anti Rat CD1d antibody, clone WTH1 efficiently blocks rat and mouse invariant natural killer T cell (iNKT cell) response to a-GalCer (an antigen for			
	iNKT cells) and has a slightly higher affinity to native CD1d than clone WTH2.			
	Clone WTH1 cross reacts weakly with mouse tissue in IHC and western blotting, but reacts strongly with both rat and mouse antigens in flow cytometry and immunoprecipitation.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.			
References	Monzon-Casanova, E. <i>et al.</i> (2010) CD1d expression in paneth cells and rat exocrine pancreas revealed by novel monoclonal antibodies which differentially affect NKT cell activation. PLoS One 5:e13089 Fichtner AS <i>et al.</i> (2015) Function and expression of CD1d and invariant natural killer			

T-cell receptor in the cotton rat (*Sigmodon hispidus*). <u>Immunology. 146 (4): 618-29.</u>
3. Monzon-Casanova, E. *et al.* (2013) Direct identification of rat iNKT cells reveals remarkable similarities to human iNKT cells and a profound deficiency in LEW rats. <u>Eur J Immunol. 43 (2): 404-15.</u>

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) <u>HRP</u>

Rabbit Anti Mouse IgG (STAR12...) RPE

Rabbit Anti Mouse IgG (STAR8...) <u>DyLight®800</u>

Goat Anti Mouse IgG (STAR76...)

RPE
Human Anti Mouse IgG2a (HCA037...)

HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Rabbit Anti Mouse IgG (STAR13...)

Goat Anti Mouse IgG (STAR70...)

Rabbit Anti Mouse IgG (STAR9...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

<u>DyLight®650</u>, <u>DyLight®680</u>, <u>DyLight®800</u>,

FITC, HRP

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA1210)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M389620:210806'

Printed on 21 Mar 2022