

Datasheet: MCA5891GA

Description:	RAT ANTI MOUSE CD283
Specificity:	CD283
Other names:	TLR3
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	11F8
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/200
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide.

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Purified murine TLR3 extracellular domain protein.
External Database Links	<p>UniProt: Q99MB1 Related reagents</p> <p>Entrez Gene: 142980 Tlr3 Related reagents</p>
Specificity	<p>Rat anti Mouse CD283 antibody, clone 11F8 recognizes mouse TLR3, otherwise known as CD283 or Toll-like receptor 3. CD283 is a 905 amino acid ~105 kDa single pass type 1 transmembrane glycoprotein associated with endosomal membranes and member of the Toll-like receptor (TLR) family. CD283 functions as a receptor for double-stranded RNA (UniProt: Q99MB1).</p> <p>TLRs are expressed on the cell surface and the endocytic compartment and recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents. TLRs also initiate cell signaling to induce production of cytokines necessary for the innate immunity and subsequent adaptive immunity.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. McCartney, S. <i>et al.</i> (2009) Distinct and complementary functions of MDA5 and TLR3 in poly(I:C)-mediated activation of mouse NK cells. J Exp Med. 206: 2967-76. 2. Jelinek, I. <i>et al.</i> (2011) TLR3-specific double-stranded RNA oligonucleotide adjuvants induce dendritic cell cross-presentation, CTL responses, and antiviral protection. J Immunol. 186: 2422-9. 3. McCartney, S.A. <i>et al.</i> (2011) RNA sensor-induced type I IFN prevents diabetes caused by a β cell-tropic virus in mice J Clin Invest. 121: 1497-507. 4. Azuma, M. <i>et al.</i> (2012) Cross-priming for antitumor CTL induced by soluble Ag + polyI:C depends on the TICAM-1 pathway in mouse CD11c(+)/CD8α(+) dendritic cells. Oncoimmunology. 1: 581-592. 5. Zeng, R. <i>et al.</i> (2016) Generation and transcriptional programming of intestinal dendritic cells: essential role of retinoic acid. Mucosal Immunol. 9 (1): 183-93. 6. Njoroge, J.M. <i>et al.</i> (2018) A flow cytometric analysis of macrophage- nanoparticle interactions <i>in vitro</i>: induction of altered Toll-like receptor expression. Int J Nanomedicine. 13: 8365-78.
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA5891GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR73...)	RPE
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight@550 , DyLight@650 , DyLight@800
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight@800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

Recommended Useful Reagents

[LEUCOPERM \(BUF09\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M404335:220822'

Printed on 21 Sep 2023