

## Datasheet: MCA5916F

<b>Description:</b>	MOUSE ANTI DOG CD25:FITC
<b>Specificity:</b>	CD25
<b>Other names:</b>	IL-2R ALPHA CHAIN
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	P4A10
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Dog		
<b>Product Form</b>	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	FITC	490	525
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
<b>Buffer Solution</b>	Phosphate buffered saline		
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin		
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml		

<b>Immunogen</b>	Enriched and stimulated canine T cells.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">O62802</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">403870</a>    IL2RA    <a href="#">Related reagents</a></p>
<b>Fusion Partners</b>	Spleen cells from immunized RBF/DnJ mice were fused with cells of the mouse P3-653 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Dog CD25, clone P4A10</b> recognizes the canine homologue of the human CD25 cell surface antigen, also known as IL-2R<math>\alpha</math>, a glycoprotein of approximately 55 kDa expressed primarily by activated T lymphocytes (<a href="#">Abrams et al. 2010</a>).</p> <p>The IL-2 receptor is composed of 3 subunits, an <math>\alpha</math> chain (CD25), a <math>\beta</math> chain (CD122) and a <math>\gamma</math> chain (CD132), CD25 functions as a low affinity receptor for IL-2.</p> <p>Antibodies to CD4,</p> <p>Mouse anti Human CD25 antibody, clone ACT1, which has demonstrated cross reactivity to canine CD25 (<a href="#">Risetto et al. 2010</a>). However this clone has poor affinity for canine CD25. The development of clone P4A10 offers a specific monoclonal Mouse anti Canine CD25, demonstrating a greater affinity for canine CD25 than the cross reactive anti human antibody (<a href="#">Abrams et al. 2010</a>).</p> <p>The dog is an important veterinary species in its own right. In addition dogs are used as an animal model in the study of a number of serious human disease states including various forms cancers (<a href="#">Paoloni et al. 2008</a>), and in genetically related diseases of the hemopoietic system (<a href="#">Bauer Jr. et al. 2009</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul
<b>References</b>	<ol style="list-style-type: none"> <li>1. Abrams, V.K. <i>et al.</i> (2010) A novel monoclonal antibody specific for canine CD25 (P4A10): selection and evaluation of canine Tregs. <a href="#">Vet Immunol Immunopathol. 135 (3-4): 257-65.</a></li> <li>2. Finn, J.D. <i>et al.</i> (2010) Eradication of neutralizing antibodies to factor VIII in canine hemophilia A after liver gene therapy. <a href="#">Blood.116: 5842-8.</a></li> <li>3. Mizutani, N. <i>et al.</i> (2020) Measurement of the concentration of serum soluble interleukin-2 receptor alpha chain in dogs with lymphoma. <a href="#">Vet Immunol Immunopathol. 225: 110054.</a></li> <li>4. Wesolowski, M. <i>et al.</i> (2023) Long-term changes of Th17 and regulatory T cells in peripheral blood of dogs with spinal cord injury after intervertebral disc herniation. <a href="#">BMC Vet Res. 19 (1): 90.</a></li> <li>5. Sheng, R. <i>et al.</i> (2023) Prognostic significance of CD25 expression in dogs with a noninvasive diagnosis of B-cell lymphoma treated with CHOP chemotherapy. <a href="#">Vet Comp Oncol. 21 (1): 28-35.</a></li> </ol>

**Further Reading**

1. Paoloni, M. & Khanna, C. (2008) Translation of new cancer treatments from pet dogs to humans. [Nat Rev Cancer. 8 \(2\): 147-56.](#)
2. Bauer, T.R. Jr *et al.* (2009) Potential large animal models for gene therapy of human genetic diseases of immune and blood cell systems. [ILAR J. 50 \(2\): 168-86.](#)
3. Risetto, K.C. *et al.* (2010) Cloning and expression of canine CD25 for validation of an anti-human CD25 antibody to compare T regulatory lymphocytes in healthy dogs and dogs with osteosarcoma. [Vet Immunol Immunopathol. 135 \(1-2\): 137-45.](#)
4. Pinheiro, D. *et al.* (2011) Phenotypic and functional characterization of a CD4(+) CD25(high) FOXP3(high) regulatory T-cell population in the dog. [Immunology. 132 \(1\): 111-22.](#)

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

**Guarantee** 12 months from date of despatch

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA5916F>  
10041

**Regulatory** For research purposes only

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

### Recommended Useful Reagents

[RAT ANTI DOG CD8 \(MCA1039GA\)](#)

[RAT ANTI DOG CD4 \(MCA1038GA\)](#)

[MOUSE ANTI DOG CD3 \(MCA1774GA\)](#)

[MOUSE ANTI DOG CD3:FITC \(MCA1774F\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M384827:210513'

Printed on 05 Mar 2024