

Datasheet: MCA5916GA

BATCH NUMBER 165730

Description:	MOUSE ANTI DOG CD25
Specificity:	CD25
Other names:	IL-2R ALPHA CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	P4A10
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen		▪		
Immunohistology - Paraffin		▪		
Immunohistology - Resin		▪		
ELISA		▪		
Immunoprecipitation		▪		
Western Blotting (1)	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)Clone P4A10 produces a double band at approximately 55kDa under non-reducing conditions. A non-specific band may be observed at 116kDa

Target Species	Dog
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Enriched and stimulated canine T cells.
External Database Links	<p>UniProt: O62802 Related reagents</p> <p>Entrez Gene: 403870 IL2RA Related reagents</p>
Fusion Partners	Spleen cells from immunized RBF/DnJ mice were fused with cells of the mouse P3-653 myeloma cell line.
Specificity	<p>Mouse anti Dog CD25, clone P4A10 recognizes the canine homologue of the human CD25 cell surface antigen, also known as IL-2Rα, a glycoprotein of approximately 55 kDa expressed primarily by activated T lymphocytes (Abrams <i>et al.</i> 2010).</p> <p>The IL-2 receptor is composed of 3 subunits, an α chain (CD25), a β chain (CD122) and a γ chain (CD132), CD25 functions as a low affinity receptor for IL-2.</p> <p>Antibodies to CD4,</p> <p>Mouse anti Human CD25 antibody, clone ACT1, which has demonstrated cross reactivity to canine CD25 (Rissetto <i>et al.</i> 2010). However this clone has poor affinity for canine CD25. The development of clone P4A10 offers a specific monoclonal Mouse anti Canine CD25, demonstrating a greater affinity for canine CD25 than the cross reactive anti human antibody (Abrams <i>et al.</i> 2010).</p> <p>The dog is an important veterinary species in its own right. In addition dogs are used as an animal model in the study of a number of serious human disease states including various forms cancers (Paoloni <i>et al.</i> 2008), and in genetically related diseases of the hemopoietic system (Bauer Jr.<i>et al.</i> 2009).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
Western Blotting	Clone P4A10 detects a double band of approximately 55kDa in PMA-IO activated canine PBMC lysates.
References	<ol style="list-style-type: none"> 1. Abrams, V.K. <i>et al.</i> (2010) A novel monoclonal antibody specific for canine CD25 (P4A10): selection and evaluation of canine Tregs. Vet Immunol Immunopathol. 135 (3-4): 257-65. 2. Finn, J.D. <i>et al.</i> (2010) Eradication of neutralizing antibodies to factor VIII in canine

hemophilia A after liver gene therapy. [Blood.116: 5842-8.](#)

3. Mizutani, N. *et al.* (2020) Measurement of the concentration of serum soluble interleukin-2 receptor alpha chain in dogs with lymphoma. [Vet Immunol Immunopathol. 225: 110054.](#)

4. Wesolowski, M. *et al.* (2023) Long-term changes of Th17 and regulatory T cells in peripheral blood of dogs with spinal cord injury after intervertebral disc herniation. [BMC Vet Res. 19 \(1\): 90.](#)

Further Reading

1. Paoloni, M. & Khanna, C. (2008) Translation of new cancer treatments from pet dogs to humans. [Nat Rev Cancer. 8 \(2\): 147-56.](#)

2. Bauer, T.R. Jr *et al.* (2009) Potential large animal models for gene therapy of human genetic diseases of immune and blood cell systems. [ILAR J. 50 \(2\): 168-86.](#)

3. Risetto, K.C. *et al.* (2010) Cloning and expression of canine CD25 for validation of an anti-human CD25 antibody to compare T regulatory lymphocytes in healthy dogs and dogs with osteosarcoma. [Vet Immunol Immunopathol. 135 \(1-2\): 137-45.](#)

4. Pinheiro, D. *et al.* (2011) Phenotypic and functional characterization of a CD4(+) CD25(high) FOXP3(high) regulatory T-cell population in the dog. [Immunology. 132 \(1\): 111-22.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA5916GA>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)

Goat Anti Mouse IgG (STAR76...) [RPE](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Goat Anti Mouse IgG (STAR70...) [FITC](#)

Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

Recommended Useful Reagents

[RAT ANTI DOG CD8 \(MCA1039GA\)](#)

[RAT ANTI DOG CD4 \(MCA1038GA\)](#)

[MOUSE ANTI DOG CD3 \(MCA1774GA\)](#)

[MOUSE ANTI DOG CD3:FITC \(MCA1774F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M381331:210512'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)