

## Datasheet: MCA5919F

<b>Description:</b>	MOUSE ANTI BOVINE CD32:FITC
<b>Specificity:</b>	CD32
<b>Other names:</b>	FcRII
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	CCG39
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10
Immunofluorescence			■	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Bovine		
<b>Species Cross Reactivity</b>	Does not react with: Sheep		
<b>Product Form</b>	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	FITC	490	525
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
<b>Buffer Solution</b>	Phosphate buffered saline		
<b>Preservative</b>	0.09% Sodium Azide (NaN <sub>3</sub> )		
<b>Stabilisers</b>	1% Bovine Serum Albumin		
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml		
<b>Immunogen</b>	Bovine FcγRII-transfected COS7 cells.		

**External Database  
Links**

**UniProt:**

[Q28110](#)   [Related reagents](#)

**Entrez Gene:**

[282229](#)   FCGR2   [Related reagents](#)

---

**Fusion Partners**   Spleen cells from immunised BALB/c mice were fused with cells of the NS-1myeloma cell line.

---

**Specificity**   **Mouse anti Bovine CD32 antibody, clone CCG39**, recognizes the bovine homologue of human CD32, one of a group of Fc receptors belonging to the immunoglobulin superfamily and involved in phagocytosis of opsonized microbes. Bovine CD32 is a single pass type 1 membrane protein of approximately 32 kDa, expressed on the cell surface of most cells including B-lymphocytes, monocytes, neutrophils and afferent veiled lymph dendritic cells ([Chattha \*et al.\* 2010](#)). It has been shown that expression of bovine CD32 is higher on macrophages than on neutrophils.

CD32 can function in an inhibitory capacity to antibody production and is the low affinity Fc receptor for IgG (FcRII), binding to the Fc region of immunoglobulin gamma [Chattha \*et al.\* 2009](#).

Mouse anti Bovine CD32, clone CCG39 is one of a number of anti bovine CD32 reagents available from Bio-Rad, clone CCG39 is of interest in that it is specific to bovine CD32 and does not recognizes ovine CD32. For detection of ovine CD32, clone [CCG36](#), which cross reacts with ovine CD32, may be used.

---

**Flow Cytometry**   Use 10ul of the suggested working dilution to label  $1 \times 10^6$  cells in 100ul

---

**Further Reading**

1. Kacskovics, I. (2004) Fc receptors in livestock species. [Vet Immunol Immunopathol. 102 \(4\): 351-62.](#)
2. Chattha, K.S. *et al.* (2009) Age related variation in expression of CD21 and CD32 on bovine lymphocytes: a cross-sectional study. [Vet Immunol Immunopathol. 130 \(1-2\): 70-8.](#)
3. Chattha, K.S. *et al.* (2010) Immunohistochemical investigation of cells expressing CD21, membrane IgM, CD32 and a follicular dendritic cell marker in the lymphoid tissues of neonatal calves. [Vet Immunol Immunopathol. 137 \(3-4\): 284-90.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.  
Storage in frost-free freezers is not recommended.  
This product should be stored undiluted. This product is photosensitive and should be protected from light.  
Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**   12 months from date of despatch

---

**Health And Safety Information**   Material Safety Datasheet documentation #10041 available at:  
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

---

**Regulatory**   For research purposes only

## Related Products

### Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:FITC \(MCA929F\)](#)

## Recommended Useful Reagents

[MOUSE ANTI BOVINE CD32:RPE \(MCA5918PE\)](#)

[MOUSE ANTI BOVINE CD32 \(MCA5918GA\)](#)

[MOUSE ANTI BOVINE CD32:FITC \(MCA5918F\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M368473:200529'

**Printed on 11 Aug 2020**

---

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)