

Datasheet: MCA5922

Description:	MOUSE ANTI HUMAN SOMATOSTATIN RECEPTOR 4
Specificity:	SOMATOSTATIN RECEPTOR 4
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	sstr4
Isotype:	IgG2a
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting		▪		

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Antibody purified from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein	IgG concentration 1mg/ml

Concentrations

Immunogen	Synthetic peptide, sequence CQQEALQPEPGRKRIPLT.
-----------	-------------------------------------------------

External Database Links

UniProt:

[P31391](#)

[Related reagents](#)

Entrez Gene:

[6754](#)

SSTR4

[Related reagents](#)

Specificity

Mouse anti Human Somatostatin Receptor 4 antibody, clone sstr4 recognizes human somatostatin receptor 4, a 388 amino acid multi pass transmembrane glycoprotein receptor for somatostatin-14. Somatostatin (SST) is a peptide hormone with wide ranging inhibitory effects on hormone secretion and cell proliferation. It is produced as a large preproSST precursor molecule that is cleaved to yield one of two active forms of the peptide: a 14 amino acid peptide (SST-14), or a 28 amino acid peptide (SST-28). The inhibitory effects of SST on hormone release and cell growth have made them candidates for the treatment of cancer and neuroendocrine disorders. Natural SSTs have very short half-lives, which led to the development of more stable synthetic derivatives, such as [octreotide](#) and [lanreotide](#), both in common clinical use.

The effects of SST are mediated via five distinct SST receptors (SSTRs). The receptors have similar affinities for natural SST-14 and SST-28, but there are marked differences in affinities towards the synthetic analogues. Somatostatin receptors are expressed by various tissues, notably neuronal, endocrine, gastrointestinal and immune cells, as well as certain tumours, with tumours often expressing more than one subtype in different combinations.

Mouse anti Human Somatostatin Receptor 4 antibody, clone sstr4 specifically recognises SSTR type 4, but none of the other SSTR subtypes.

Histology Positive Control Tissue

Human lung tissue

References

1. Leijon, H. *et al.* (2019) Variable somatostatin receptor subtype expression in 151 primary pheochromocytomas and paragangliomas. [Hum Pathol. 86: 66-75.](#)
2. Ozkaya, H.M. *et al.* (2018) Germline mutations of aryl hydrocarbon receptor-interacting protein (AIP) gene and somatostatin receptor 1-5 and AIP immunostaining in patients with sporadic acromegaly with poor versus good response to somatostatin analogues. [Pituitary. 21 \(4\): 335-346.](#)
3. Remes, S.M. *et al.* (2019) Immunohistochemical Expression of Somatostatin Receptor Subtypes in a Panel of Neuroendocrine Neoplasias. [J Histochem Cytochem. 67 \(10\): 735-43.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA5922 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	------------------	-------------------------------------------------------------------------------------------------------------------------------------------------	---------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M429985:240501'

Printed on 01 May 2024