Datasheet: MCA5934 BATCH NUMBER 164173

Description:	RAT ANTI MOUSE JAM-C
Specificity:	JAM-C
Other names:	JAM-3
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CRAM-19 H36
Isotype:	lgG2a
Quantity:	0.25 mg

Product Details

ApplicationsThis product has been reported to work in the following applications. This information is
derived from testing within our laboratories, peer-reviewed publications or personal
communications from the originators. Please refer to references indicated for further
information. For general protocol recommendations, please visit www.bio-
rad-antibodies.com/protocols.

		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry				1/50 - 1/100
	Immunohistology - Frozen	•			
	Immunohistology - Paraffin			•	
	ELISA				
	Immunoprecipitation	•			
	Western Blotting			-	
	Immunofluorescence				
	Where this product has n	ot been t	ested for	use in a particular tech	nique this does not
	•			•	•
	necessarily exclude its us	se in suicr			
	necessarily exclude its us		•		
	a guide only. It is recomm		•		
	•	nended th	nat the use	er titrates the product f	
Target Species	a guide only. It is recomm system using appropriate	nended th	nat the use	er titrates the product f	
Target Species	a guide only. It is recomm	nended th	nat the use	er titrates the product f	
	a guide only. It is recomm system using appropriate	nended th	nat the use	er titrates the product f	
Species Cross	a guide only. It is recomm system using appropriate Mouse Reacts with: Human	nended th negative	hat the use	er titrates the product t	for use in their own
Species Cross	a guide only. It is recomm system using appropriate Mouse Reacts with: Human N.B. Antibody reactivity a	nended th negative nd worki	nat the use positive of ng conditi	er titrates the product f controls.	for use in their own
Species Cross	a guide only. It is recommon system using appropriate Mouse Reacts with: Human N.B. Antibody reactivity a reactivity is derived from	nended th negative nd worki testing w	nat the use positive ng conditi ithin our la	er titrates the product t controls. ons may vary betweer aboratories, peer-revie	for use in their own n species. Cross wed publications or
Target Species Species Cross Reactivity	a guide only. It is recommon system using appropriate Mouse Reacts with: Human N.B. Antibody reactivity a reactivity is derived from the personal communications	nended th negative nd worki testing w	nat the use positive ng conditi ithin our la	er titrates the product t controls. ons may vary betweer aboratories, peer-revie	for use in their own n species. Cross wed publications or
Species Cross	a guide only. It is recommon system using appropriate Mouse Reacts with: Human N.B. Antibody reactivity a reactivity is derived from	nended th negative nd worki testing w	nat the use positive ng conditi ithin our la	er titrates the product t controls. ons may vary betweer aboratories, peer-revie	for use in their own n species. Cross wed publications or

Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant			
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)			
Carrier Free	Yes			
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml			
Immunogen	Recombinant soluble JAM-C.			
External Database Links	UniProt:			
	Q9D8B7 Related reagents			
	Q9BX67 Related reagents			
	Entrez Gene:			
	83964 Jam3 Related reagents			
	83700 JAM3 Related reagents			
Fusion Partners	Spleen cells from immunised Fischer rats were fused with cells of the Sp2/0 myeloma cell line.			
Specificity	Rat anti Mouse JAM-C, clone CRAM-19 H36 recognizes mouse and human Junctional adhesion molecule C (JAM-C), also known as JAM-3 and, historically, as JAM-2.			
	JAM-C is expressed at junctions between endothelial and epithelial cells, as well as on leukocytes, platelets, vascular smooth muscle cells and fibroblasts, amongst other cell			
	types. It plays a role in tight junctions and inflammatory processes and interacts with JAM-A and JAM-B.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.			
References	 Aurrand-lions, M. <i>et al.</i> (2001) JAM-2, a novel immunoglobulin superfamily molecule, expressed by endothelial and lymphatic cells. <u>J Biol Chem. 276 (4): 2733-41.</u> Johnson-léger, C.A. <i>et al.</i> (2002) Junctional adhesion molecule-2 (JAM-2) promotes lymphocyte transendothelial migration. <u>Blood. 100 (7): 2479-86.</u> Morita, Y. <i>et al.</i> (2010) Heterogeneity and hierarchy within the most primitive hematopoietic stem cell compartment. <u>J Exp Med. 207 (6): 1173-82.</u> 			
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.			
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in			

	frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA5934 10040
Regulatory	For research purposes only

Related Products

America

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16)	DyLight®800				
Rabbit Anti Rat IgG (STAR17)	<u>FITC</u>				
Goat Anti Rat IgG (STAR72)	HRP				
Goat Anti Rat IgG (STAR69)	<u>FITC</u>				
Goat Anti Rat IgG (STAR73)	RPE				
Rabbit Anti Rat IgG (STAR21)	HRP				
Goat Anti Rat IgG (MOUSE ADSORBED)	(STAR71)DyLight®550, DyLight®650, DyLight®800				
Goat Anti Rat IgG (STAR131)	<u>Alk. Phos.</u> , <u>Biotin</u>				
Recommended Negative Controls					
RAT IgG2a NEGATIVE CONTROL (MCA1212)					
North & South Tel: +1 800 265 7376 Worldwide	e Tel: +44 (0)1865 852 700 Europe Tel: +49 (0) 89 8090 95 21				

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M381777:210512'

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com