

Datasheet: MCA5935

## **BATCH NUMBER 1112**

| Description:            | RAT ANTI MOUSE JAM-C            |
|-------------------------|---------------------------------|
| Specificity:            | JAM-C                           |
| Other names:            | JAM-3                           |
| Format:                 | Purified                        |
|                         |                                 |
| Product Type:           | Monoclonal Antibody             |
| Product Type:<br>Clone: | Monoclonal Antibody CRAM-18 F26 |
|                         | •                               |

# **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                            | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry             |     |    |                | 1/10 - 1/25        |
| Immunohistology - Frozen   |     |    |                |                    |
| Immunohistology - Paraffin |     |    |                |                    |
| ELISA                      |     |    | •              |                    |
| Immunoprecipitation        |     |    |                |                    |
| Western Blotting           |     |    |                |                    |
| Immunofluorescence         | •   |    |                |                    |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species              | Mouse   |
|-----------------------------|---|
| Species Cross<br>Reactivity | Reacts with: Human  N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. |
| Product Form                | Purified IgG - liquid   |

| Preparation                       | Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant   |  |  |
|-----------------------------------|---|--|--|
| Buffer Solution                   | Phosphate buffered saline   |  |  |
| Preservative<br>Stabilisers       | 0.09% Sodium Azide (NaN <sub>3</sub> )  |  |  |
| Carrier Free                      | Yes   |  |  |
| Approx. Protein<br>Concentrations | IgG concentration 1.0 mg/ml   |  |  |
| Immunogen                         | Recombinant soluble JAM-C.  |  |  |
| External Database<br>Links        | UniProt:  Q9D8B7 Related reagents   |  |  |
|                                   | Q9BX67 Related reagents   |  |  |
|                                   | Entrez Gene:  |  |  |
|                                   | 83964 Jam3 <u>Related reagents</u> 83700 JAM3 <u>Related reagents</u>   |  |  |
| Fusion Partners                   | Spleen cells from immunised Fischer rats were fused with cells of the Sp2/0 myeloma cell line.  |  |  |
| Specificity                       | Rat anti Mouse JAM-c antibody, clone CRAM-18 F26 recognizes mouse and human Junctional adhesion molecule C (JAM-C), also known as JAM-3 and, historically, as JAM-2.  |  |  |
|                                   | JAM-C is expressed at junctions between endothelial and epithelial cells, as well as on leukocytes, platelets, vascular smooth muscle cells and fibroblasts, amongst other cell types. It plays a role in tight junctions and inflammatory processes and interacts with JAM-A and JAM-B.  |  |  |
|                                   | Clone CRAM-18 F26 has been reported to inhibit transendothelial migration ( <u>Johnson-Léger <i>et al.</i> 2002</u> ).  |  |  |
| Flow Cytometry                    | Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.   |  |  |
| References                        | <ol> <li>Aurrand-lions, M. <i>et al.</i> (2001) Heterogeneity of endothelial junctions is reflected by differential expression and specific subcellular localization of the three JAM family members. Blood. 98 (13): 3699-707.</li> <li>Johnson-léger, C.A. <i>et al.</i> (2002) Junctional adhesion molecule-2 (JAM-2) promotes lymphocyte transendothelial migration. Blood. 100 (7): 2479-86.</li> <li>Aurrand-lions, M. <i>et al.</i> (2001) JAM-2, a novel immunoglobulin superfamily molecule, expressed by endothelial and lymphatic cells. J Biol Chem. 276 (4): 2733-41.</li> <li>Forsberg, E.C. <i>et al.</i> (2005) Differential expression of novel potential regulators in</li> </ol> |  |  |

hematopoietic stem cells. PLoS Genet. 1(3):e28.

5. Miranda, J. *et al.* (2019) Syncytiotrophoblast of Placentae from Women with Zika Virus Infection Has Altered Tight Junction Protein Expression and Increased Paracellular Permeability. Cells. 8 (10)Sep 29 [Epub ahead of print].

Storage Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend

microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:

https://www.bio-rad-antibodies.com/SDS/MCA5935

10040

**Regulatory** For research purposes only

# Related Products

# **Recommended Secondary Antibodies**

Rabbit Anti Rat IgG (STAR16...) DyLight®800

Rabbit Anti Rat IgG (STAR17...)

Goat Anti Rat IgG (STAR69...)

Goat Anti Rat IgG (STAR73...)

Rabbit Anti Rat IgG (STAR21...)

HRP

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) <u>DyLight®550</u>, <u>DyLight®650</u>, <u>DyLight®800</u>

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

Goat Anti Rat IgG (STAR72...) HRP

## **Recommended Negative Controls**

#### RAT IgG2a NEGATIVE CONTROL (MCA1212)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368492:200529'

## Printed on 29 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint