

Datasheet: MCA5935

Description:	RAT ANTI MOUSE JAM-C
Specificity:	JAM-C
Other names:	JAM-3
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CRAM-18 F26
Isotype:	IgG2a
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/25
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Species Cross Reactivity	Reacts with: Human N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein	IgG concentration 1.0 mg/ml

Concentrations

Immunogen Recombinant soluble JAM-C.

External Database Links

UniProt:

[Q9D8B7](#) [Related reagents](#)

[Q9BX67](#) [Related reagents](#)

Entrez Gene:

[83964](#) Jam3 [Related reagents](#)

[83700](#) JAM3 [Related reagents](#)

Fusion Partners Spleen cells from immunised Fischer rats were fused with cells of the Sp2/0 myeloma cell line.

Specificity **Rat anti Mouse JAM-c antibody, clone CRAM-18 F26** recognizes mouse and human Junctional adhesion molecule C (JAM-C), also known as JAM-3 and, historically, as JAM-2.

JAM-C is expressed at junctions between endothelial and epithelial cells, as well as on leukocytes, platelets, vascular smooth muscle cells and fibroblasts, amongst other cell types. It plays a role in tight junctions and inflammatory processes and interacts with JAM-A and JAM-B.

Clone CRAM-18 F26 has been reported to inhibit transendothelial migration ([Johnson-Léger *et al.* 2002](#)).

Flow Cytometry Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References

1. Aurrand-lions, M. *et al.* (2001) Heterogeneity of endothelial junctions is reflected by differential expression and specific subcellular localization of the three JAM family members. [Blood. 98 \(13\): 3699-707.](#)
 2. Johnson-léger, C.A. *et al.* (2002) Junctional adhesion molecule-2 (JAM-2) promotes lymphocyte transendothelial migration. [Blood. 100 \(7\): 2479-86.](#)
 3. Aurrand-lions, M. *et al.* (2001) JAM-2, a novel immunoglobulin superfamily molecule, expressed by endothelial and lymphatic cells. [J Biol Chem. 276 \(4\): 2733-41.](#)
 4. Forsberg, E.C. *et al.* (2005) Differential expression of novel potential regulators in hematopoietic stem cells. [PLoS Genet. 1\(3\):e28.](#)
 5. Miranda, J. *et al.* (2019) Syncytiotrophoblast of Placentae from Women with Zika Virus Infection Has Altered Tight Junction Protein Expression and Increased Paracellular Permeability. [Cells. 8 \(10\)Sep 29 \[Epub ahead of print\].](#)
-

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (STAR72...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®800

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	------------------	-------------------------------------------------------------------------------------------------------------------------------------------------	---------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

'M368492:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)