

Datasheet: MCA5935T

BATCH NUMBER 169064

Description:	RAT ANTI MOUSE JAM-C
Specificity:	JAM-C
Other names:	JAM-3
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CRAM-18 F26
Isotype:	IgG2a
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/25
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Species Cross Reactivity	<p>Reacts with: Human</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant soluble JAM-C.
External Database Links	<p>UniProt:</p> <p>Q9D8B7 Related reagents</p> <p>Q9BX67 Related reagents</p> <p>Entrez Gene:</p> <p>83964 Jam3 Related reagents</p> <p>83700 JAM3 Related reagents</p>
Fusion Partners	Spleen cells from immunized Fischer rats were fused with cells of the Sp2/0 myeloma cell line.
Specificity	<p>Rat anti Mouse JAM-c antibody, clone CRAM-18 F26 recognizes mouse and human Junctional adhesion molecule C (JAM-C), also known as JAM-3 and, historically, as JAM-2.</p> <p>JAM-C is expressed at junctions between endothelial and epithelial cells, as well as on leukocytes, platelets, vascular smooth muscle cells and fibroblasts, amongst other cell types. It plays a role in tight junctions and inflammatory processes and interacts with JAM-A and JAM-B.</p> <p>Rat anti Mouse JAM-c antibody, clone CRAM-18 F26 has been reported to inhibit transendothelial migration (Johnson-Léger <i>et al.</i> 2002).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Aurrand-lions, M. <i>et al.</i> (2001) JAM-2, a novel immunoglobulin superfamily molecule, expressed by endothelial and lymphatic cells. J Biol Chem. 276 (4): 2733-41. 2. Aurrand-lions, M. <i>et al.</i> (2001) Heterogeneity of endothelial junctions is reflected by differential expression and specific subcellular localization of the three JAM family members. Blood. 98 (13): 3699-707. 3. Johnson-léger, C.A. <i>et al.</i> (2002) Junctional adhesion molecule-2 (JAM-2) promotes lymphocyte transendothelial migration. Blood. 100 (7): 2479-86. 4. Forsberg, E.C. <i>et al.</i> (2005) Differential expression of novel potential regulators in

hematopoietic stem cells. [PLoS Genet. 1\(3\):e28.](#)

5. Radulovic, V. *et al.* (2019) Junctional Adhesion Molecule 2 Represents a Subset of Hematopoietic Stem Cells with Enhanced Potential for T Lymphopoiesis. [Cell Rep. 27\(10\): 2826-2836.e5.](#)

6. Miranda, J. *et al.* (2019) Syncytiotrophoblast of Placentae from Women with Zika Virus Infection Has Altered Tight Junction Protein Expression and Increased Paracellular Permeability. [Cells. 8 \(10\)Sep 29 \[Epub ahead of print\].](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA5935T>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)

[DyLight®800](#)

Rabbit Anti Rat IgG (STAR17...)

[FITC](#)

Goat Anti Rat IgG (STAR69...)

[FITC](#)

Goat Anti Rat IgG (STAR73...)

[RPE](#)

Rabbit Anti Rat IgG (STAR21...)

[HRP](#)

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) [DyLight®550](#), [DyLight®650](#), [DyLight®800](#)

Goat Anti Rat IgG (STAR131...)

[Alk. Phos.](#), [Biotin](#)

Goat Anti Rat IgG (STAR72...)

[HRP](#)

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M382420:210513'

Printed on 19 Dec 2024