

## Datasheet: MCA594PE

<b>Description:</b>	MOUSE ANTI HUMAN CD42a:RPE
<b>Specificity:</b>	CD42a
<b>Other names:</b>	GPIX
<b>Format:</b>	RPE
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	FMC-25
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human		
<b>Product Form</b>	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
<b>Reconstitution</b>	Reconstitute with 1.0 ml distilled water		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	RPE 488nm laser	496	578
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		

<b>Buffer Solution</b>	Phosphate buffered saline.
<b>Preservative</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Stabilisers</b>	1% Bovine Serum Albumin
<b>Immunogen</b>	Peripheral blood mononuclear cells.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P14770</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">2815</a>    GP9    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_10964918
<b>Specificity</b>	<p><b>Mouse anti Human CD42a antibody, clone FMC-25</b> recognizes human CD42a, also known as Platelet glycoprotein IX, Glycoprotein 9 or GP-IX. CD42a is a 177 amino acid, ~20kDa type I single pass transmembrane glycoprotein containing a single <a href="#">leucine-rich repeat containing N-terminal</a> domain and a single <a href="#">leucine-rich repeat containing C-terminal</a> domain.</p> <p>CD42a is expressed by platelets and megakaryocytes and forms a covalent complex with CD42c (GP-1b-beta), CD42b (GP-1b-alpha) and CD42d (platelet glycoprotein V) to create the platelet surface receptor for von Willebrand factor. Incubation of the intact von Willebrand receptor complex with clone FMC-25 does not appear to inhibit binding of von Willebrand factor to the receptor (<a href="#">Yan et al. 2011</a>). Defects in the GP1BB gene encoding human CD42a can lead to the inherited bleeding disorder Bernard-Soulier syndrome (<a href="#">Diz-Küçükkaya 2013</a>), characterized by prolonged bleeding times, thrombocytopenia and the appearance of giant platelets in the circulation (<a href="#">Johns et al. 2014</a>).</p> <p>Mouse anti human CD42a antibody, clone FMC-25 has been successfully used as a capture reagent for platelet-autoantibody complexes in the sera of patients presenting thrombocytopenia associated with antiphospholipid syndrome (<a href="#">Godeau et al. 1997</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 1 x 10 <sup>6</sup> cells in 100ul.
<b>Histology Positive Control Tissue</b>	Bone marrow
<b>References</b>	<ol style="list-style-type: none"> <li>Zola, H. <i>et al.</i> (1984) Monoclonal antibodies against antigens of the human platelet surface: preparation and properties. <a href="#">Pathology. 16 (1): 73-8.</a></li> <li>Berndt, M.C. <i>et al.</i> (1985) Molecular characterization of quinine/quinidine drug-dependent antibody platelet interaction using monoclonal antibodies. <a href="#">Blood. 66 (6): 1292-301.</a></li> <li>Berndt, M.C. <i>et al.</i> (1985) Purification and preliminary characterization of the glycoprotein Ib complex in the human platelet membrane. <a href="#">Eur J Biochem. 151 (3): 637-49.</a></li> <li>Berndt, M.C. <i>et al.</i> (1983) Additional glycoprotein defects in Bernard-Soulier's</li> </ol>

- syndrome: confirmation of genetic basis by parental analysis. [Blood. 62 \(4\): 800-7.](#)
5. San Miguel, J.F. *et al.* (1985) Characterization of blast cells in chronic granulocytic leukaemia in transformation, acute myelofibrosis and undifferentiated leukaemia. II. Studies with monoclonal antibodies and terminal transferase. [Br J Haematol. 59 \(2\): 297-309.](#)
  6. San Miguel, J.F. *et al.* (1986) Surface marker analysis in acute myeloid leukaemia and correlation with FAB classification. [Br J Haematol. 64 \(3\): 547-60.](#)
  7. Smith GA *et al.* (2007) Severe fetomaternal alloimmune thrombocytopenia due to anti-human platelet antigen (HPA)-1a in a mother with a rare and silenced ITGB3\*0101 (GPIIIa) allele. [Vox Sang. 93 \(4\): 325-30.](#)
  8. Berndt, M.C. *et al.* (1988) Ristocetin-dependent reconstitution of binding of von Willebrand factor to purified human platelet membrane glycoprotein Ib-IX complex. [Biochemistry. 27 \(2\): 633-40.](#)
  9. Yan, R. *et al.* (2011) Reconstitution of the platelet glycoprotein Ib-IX complex in phospholipid bilayer Nanodiscs. [Biochemistry. 50: 10598-606.](#)
  10. Sailer, T. *et al.* (2006) The course of severe autoimmune thrombocytopenia in patients not undergoing splenectomy. [Haematologica. 91: 1041-5.](#)
  11. Tomicic, M. *et al.* (2006) Frequency of HPA-15a and HPA-15b (Gov a/b) human platelet alloantigens in the Croatian population. [Arch Med Res. 37: 172-4.](#)
  12. Starcevic, M. *et al.* (2010) Neonatal alloimmune thrombocytopenia caused by anti-HLA-A24 alloantibodies. [Acta Paediatr. 99: 630-2.](#)
  13. Schallmoser, K. *et al.* (2006) Specificities of platelet autoantibodies and platelet activation in lupus anticoagulant patients: a relation to their history of thromboembolic disease. [Lupus. 15: 507-14.](#)
  14. Meyer, O. *et al.* (2003) Diclofenac-induced antibodies against RBCs and platelets: two case reports and a concise review. [Transfusion. 43: 345-9.](#)
  15. Lubenow, N. *et al.* (2000) Very low platelet counts in post-transfusion purpura falsely diagnosed as heparin-induced thrombocytopenia. Report of four cases and review of literature. [Thromb Res. 100: 115-25.](#)
  16. Ghevaert, C. *et al.* (2008) A nonsynonymous SNP in the ITGB3 gene disrupts the conserved membrane-proximal cytoplasmic salt bridge in the alphaIIb beta3 integrin and cosegregates dominantly with abnormal proplatelet formation and macrothrombocytopenia. [Blood. 111: 3407-14.](#)
  17. Bub, C.B. *et al.* (2016) The use of a potential novel tool in virtual crossmatching for platelet transfusion in platelet refractoriness. [Vox Sang. 110 \(1\): 70-8.](#)
  18. Michel, M. *et al.* (2002) Platelet autoantibodies and lupus-associated thrombocytopenia. [Br J Haematol. 119 \(2\): 354-8.](#)
  19. Schallmoser, K. *et al.* (2006) Delayed detectability of anti-HPA-3a by the MAIPA assay in a severe neonatal alloimmune thrombocytopenia, but successful transfusion of incompatible donor platelets: a case report. [Vox Sang. 91 \(2\): 181-3.](#)
  20. Wihadmadyatami, H. *et al.* (2015) Alloantibody against new platelet alloantigen (Lap(a)) on glycoprotein IIb is responsible for a case of fetal and neonatal alloimmune thrombocytopenia. [Transfusion. 55 \(12\): 2920-9.](#)

---

**Storage**

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend

microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of reconstitution.
------------------	--

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10041 available at: 10041: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

From March 15, 2021, we will no longer supply printed datasheets with our products.  
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](https://www.bio-rad-antibodies.com)

'M335044:181203'

**Printed on 10 Feb 2021**