Datasheet: MCA5950GA BATCH NUMBER 161064

Description:	MOUSE ANTI HUMAN CD44v6		
Specificity:	CD44v6		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	VFF-18		
lsotype:	lgG1		
Quantity:	0.1 mg		

Product Details

Applications	This product has been re derived from testing withi communications from the information. For general p rad-antibodies.com/proto	peer-reviewed publica e refer to references ir	ces indicated for further			
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry	-				
	Immunohistology - Frozen	-			1.0 - 2.0ug/ml	
	Immunohistology - Paraffin (1)	-			1.0 - 2.0ug/ml	
	ELISA	-				
	Immunoprecipitation					
	Western Blotting	-				
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. (1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Tris/EDTA buffer pH 9.0 or Sodium citrate buffer pH 6.0 is recommended for this purpose.					
Target Species	Human					
Product Form	Purified IgG - liquid					
Preparation	Purified IgG prepared by affinity chromatography on Protein G					
Buffer Solution	Phosphate buffered saline					
Preservative	<0.1% Sodium Azide (NaN ₃)					

Stabilisers		
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml	
Immunogen	Glutathione S Transferase (GST) fusion protein corresponding to the variable domain v6 of human CD44.	
External Database Links	UniProt: <u>P16070</u> <u>Related reagents</u>	
	Entrez Gene: <u>960</u> CD44 <u>Related reagents</u>	
Synonyms	LHR, MDU2, MDU3, MIC4	
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the P3X63Ag8.653 myeloma cell line.	
Specificity Mouse anti Human CD44v6, clone VFF-18 recognizes a centrally located 14mer encoded by amino acid residues 18-31, within variant exon 6 (v6) of human CD44 (<u>Reinisch <i>et al</i>, 1998</u>). CD44 is a member of a polymorphic family of transmembra glycoproteins, which exists as a broadly expressed ~80-90 kDa standard form (CD and also as numerous splice variants, due to extensive alternative splicing and post-translational modifications.		
	CD44s is involved in cell adhesion, lymphocyte activation, signal transduction, haematopoiesis, recirculation and homing, and CD44v6 mediates leukocyte to endothelial cell attachment and rolling, as well as homing to sites of inflammation and peripheral organs.	
	As well as its role as a receptor for hyaluronic acid (HA), CD44s also interacts with other ligands including matrix metalloproteinases (MMPs), fibronectin and collagens, and has been linked with cancer of the breast, colon, stomach and squamous cell carcinomas (HNSCC) of the head and neck, as well as being a prognostic marker for non-Hodgkin's lymphoma.	
	The variant isoforms of CD44 (CD44v) are highly expressed on epithelial cells but not leukocytes, and their association with cell transformation and cancer is also well established, supported by numerous immunohistochemical investigations including those looking at HNSCC. Mouse anti Human CD44v6, clone VFF-18 binds to CD44v6 with high affinity when compared with anti-CD44v6 clone 2F10, and is suitable for detecting CD44v6 in inflammatory bowel disease (IBD) (<u>Reinisch <i>et al.</i> 1998</u>).	
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul	
References	1. Reinisch, W. <i>et al.</i> (1998) Poor diagnostic value of colonic CD44v6 expression and serum concentrations of its soluble form in the differentiation of ulcerative colitis from	

	 Crohn's disease. <u>Gut. 43 (3): 375-82.</u> 2. Foekens, J.A. <i>et al.</i> (1999) Prognostic value of CD44 variant expression in primary breast cancer. <u>Int J Cancer. 84 (3): 209-15.</u> 3. Adolf, G.R. and Patzelt, E. (1999) Monoclonal antibody against CD44v6 <u>US patent pub No: US08750359</u> 4. Odenthal, J. <i>et al.</i> (2018) Targeting CD44v6 for fluorescence-guided surgery in head and neck squamous cell carcinoma. <u>Sci Rep. 8 (1): 10467.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in
	frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA5950GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	RPE			
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>				
Goat Anti Mouse IgG (STAR76)	RPE			
Rabbit Anti Mouse IgG (STAR13)	HRP			
Goat Anti Mouse IgG (STAR70)	FITC			
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,			
	DyLight®650, DyLight®680, DyLight®800,			
	$\underline{Dylight@000}, \underline{Dylight@000}, \underline{Dylight@000},$			
	FITC, HRP			
Rabbit Anti Mouse IgG (STAR9)				
Rabbit Anti Mouse IgG (STAR9) Goat Anti Mouse IgG (STAR77)	FITC, HRP			
3 ()	FITC, HRP FITC			
Goat Anti Mouse IgG (STAR77)	FITC, HRP FITC HRP			

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bi	o-rad.com	Email: antibody_sales_uk@bio	o-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M389632:210806'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint