



Datasheet: MCA5950GA

BATCH NUMBER 170290

Description:	MOUSE ANTI HUMAN CD44v6
Specificity:	CD44v6
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	VFF-18
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			1.0 - 2.0ug/ml
Immunohistology - Paraffin (1)	▪			1.0 - 2.0ug/ml
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Tris/EDTA buffer pH 9.0 or Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Glutathione S Transferase (GST) fusion protein corresponding to the variable domain v6 of human CD44.
External Database Links	<p>UniProt: P16070 Related reagents</p> <p>Entrez Gene: 960 CD44 Related reagents</p>
Synonyms	LHR, MDU2, MDU3, MIC4
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the P3X63Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Human CD44v6, clone VFF-18 recognizes a centrally located 14mer epitope encoded by amino acid residues 18-31, within variant exon 6 (v6) of human CD44 (Reinisch et al, 1998). CD44 is a member of a polymorphic family of transmembrane glycoproteins, which exists as a broadly expressed ~80-90 kDa standard form (CD44s), and also as numerous splice variants, due to extensive alternative splicing and post-translational modifications.</p> <p>CD44s is involved in cell adhesion, lymphocyte activation, signal transduction, haematopoiesis, recirculation and homing, and CD44v6 mediates leukocyte to endothelial cell attachment and rolling, as well as homing to sites of inflammation and peripheral organs.</p> <p>As well as its role as a receptor for hyaluronic acid (HA), CD44s also interacts with other ligands including matrix metalloproteinases (MMPs), fibronectin and collagens, and has been linked with cancer of the breast, colon, stomach and squamous cell carcinomas (HNSCC) of the head and neck, as well as being a prognostic marker for non-Hodgkin's lymphoma.</p> <p>The variant isoforms of CD44 (CD44v) are highly expressed on epithelial cells but not leukocytes, and their association with cell transformation and cancer is also well established, supported by numerous immunohistochemical investigations including those looking at HNSCC. Mouse anti Human CD44v6, clone VFF-18 binds to CD44v6 with high affinity when compared with anti-CD44v6 clone 2F10, and is suitable for detecting CD44v6 in inflammatory bowel disease (IBD) (Reinisch et al. 1998).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul
References	1. Foekens, J.A. <i>et al.</i> (1999) Prognostic value of CD44 variant expression in primary

breast cancer. [Int J Cancer. 84 \(3\): 209-15.](#)

2. Odenthal, J. *et al.* (2018) Targeting CD44v6 for fluorescence-guided surgery in head and neck squamous cell carcinoma. [Sci Rep. 8 \(1\): 10467.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA5950GA>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M410626:221028'

Printed on 02 Jun 2026