

# Datasheet: MCA5951F

**BATCH NUMBER 168269**

|                      |                         |
|----------------------|-------------------------|
| <b>Description:</b>  | MOUSE ANTI PIG CD3:FITC |
| <b>Specificity:</b>  | CD3 EPSILON             |
| <b>Format:</b>       | FITC                    |
| <b>Product Type:</b> | Monoclonal Antibody     |
| <b>Clone:</b>        | PPT3                    |
| <b>Isotype:</b>      | IgG1                    |
| <b>Quantity:</b>     | 0.1 mg                  |

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

|                | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪   |    |                | Neat - 1/10        |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

|                          |   |                     |                   |
|--------------------------|---|---------------------|-------------------|
| Target Species           | Pig   |                     |                   |
| Species Cross Reactivity | Does not react with:Bovine, Goat, Horse, Human, Sheep   |                     |                   |
| Product Form             | Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid                |                     |                   |
| Max Ex/Em                | Fluorophore   | Excitation Max (nm) | Emission Max (nm) |
|                          | FITC  | 490                 | 525               |
| Preparation              | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant |                     |                   |
| Buffer Solution          | Phosphate buffered saline   |                     |                   |
| Preservative Stabilisers | 0.09% Sodium Azide (NaN <sub>3</sub> )<br>1% Bovine Serum Albumin                             |                     |                   |

|                                |  |
|--------------------------------|--|
| Approx. Protein Concentrations | IgG concentration 0.1 mg/ml  |
| Immunogen                      | Porcine PBMCs  |
| External Database Links        | <p><b>UniProt:</b><br/> <a href="#">Q7YRN2</a>   <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b><br/> <a href="#">397455</a>   CD3E   <a href="#">Related reagents</a></p>   |
| Fusion Partners                | Lymph node cells from immunized BALB/c mice were fused with cells of the NS0 myeloma cell line   |
| Specificity                    | <p><b>Mouse anti Pig CD3, clone PPT3</b> recognizes the porcine homologue of human CD3ε, a 24 kDa single pass type I membrane protein expressed by T-lymphocytes. Clone PPT3, also known under the clone designation FY1H2, was clustered at the second international swine CD workshop and found to specifically recognise an epitope on the porcine CD3ε designated as CD3c (<a href="#">Pescovitz, M.D., et al. 1998</a>).</p> <p>CD3 is a multimeric protein complex composed of four distinct polypeptide chains (ε, γ, δ, ζ) that assemble and function as three pairs of dimers (εγ, εδ, ζζ). The CD3 complex serves as a T cell co-receptor that associates non-covalently with the T cell receptor (TCR) (<a href="#">Guy, C.S &amp; Vignali, D.G. 2009</a>). CD3 is a defining feature of cells belonging to the T cell lineage, antibodies recognising pig CD3 therefore provide useful markers of porcine T cells.</p> <p>Clone PPT3 has been demonstrated to recognise an epitope that is expressed both intracellularly and extracellularly, additionally clone PPT3 has been demonstrated to activate α/β T-cells (<a href="#">Kirkham P.A., et al. 1996</a>).</p> <p>Clone PPT3 was tested on PBL from a range of other mammalian species and found to be negative suggesting that the epitope recognised by this clone is specific to porcine (<a href="#">Yang, H. et al. 1996</a>).</p> |
| References                     | <ol style="list-style-type: none"> <li>1. Kirkham, P.A. <i>et al.</i> (1996) Porcine CD3 epsilon: its characterization, expression and involvement in activation of porcine T lymphocytes. <a href="#">Immunology. 87 (4): 616-23.</a></li> <li>2. Uehlein, S. <i>et al.</i> (2021) Human-like Response of Pig T Cells to Superagonistic Anti-CD28 Monoclonal Antibodies. <a href="#">J Immunol. 207 (10): 2473-88.</a></li> <li>3. Zhao, H. <i>et al.</i> (2022) Development of <i>RAG2</i><sup>-/-</sup> <i>IL2Rγ</i><sup>-/-</sup> immune deficient FAH-knockout miniature pig. <a href="#">Front Immunol. 13: 950194.</a></li> <li>4. Maciag, S.S. <i>et al.</i> (2022) On the influence of the source of porcine colostrum in the development of early immune ontogeny in piglets. <a href="#">Sci Rep. 12 (1): 15630.</a></li> <li>5. dos Santos, M.C. <i>et al.</i> (2023) Effect of yeast extracted β-glucans on the immune response and reproductive performance of gilts in the adaptation, gestation, and lactation periods <a href="#">Livestock Science. 275: 105289.</a></li> <li>6. Haach, V. <i>et al.</i> (2023) A polyvalent virosomal influenza vaccine induces broad cellular</li> </ol>   |

and humoral immunity in pigs. [Virol J. 20 \(1\): 181.](#)

7. Hu, Z. *et al.* (2019) Genomic variant in porcine TNFRSF1A gene and its effects on TNF signaling pathway in vitro. [Gene. 700: 105-109.](#)

---

|                        |   |
|------------------------|---|
| <b>Further Reading</b> | 1. Guy, C.S. & Vignali, D.A. (2009) Organization of proximal signal initiation at the TCR:CD3 complex. <a href="#">Immunol Rev. 32: 7-21.</a> |
|------------------------|---|

---

|                |   |
|----------------|---|
| <b>Storage</b> | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. |
|----------------|---|

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

|                  |                                 |
|------------------|---------------------------------|
| <b>Guarantee</b> | 12 months from date of despatch |
|------------------|---------------------------------|

---

|                                      |  |
|--------------------------------------|--|
| <b>Health And Safety Information</b> | Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA5951F10041">https://www.bio-rad-antibodies.com/SDS/MCA5951F10041</a> |
|--------------------------------------|--|

---

|                   |                            |
|-------------------|----------------------------|
| <b>Regulatory</b> | For research purposes only |
|-------------------|----------------------------|

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

### Recommended Useful Reagents

[MOUSE ANTI PIG CD4 ALPHA:RPE \(MCA1749PE\)](#)

[MOUSE ANTI PIG wCD8 ALPHA:RPE \(MCA1223PE\)](#)

[MOUSE ANTI PIG CD25 \(MCA1736GA\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M384683:210513'

Printed on 17 May 2024