

# Datasheet: MCA5951F BATCH NUMBER 1807

Description:	MOUSE ANTI PIG CD3:FITC
Specificity:	CD3 EPSILON
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	PPT3
lsotype:	lgG1
Quantity:	0.1 mg

# **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry	-			Neat - 1/10	
	Where this product has not been tested for use in a particular technique this does not					
	necessarily exclude it: a guide only. It is reco system using appropr	mmended tha	t the use	r titrates the product	ng dilutions are given as for use in their own	
Target Species	Pig					
Species Cross Reactivity	Does not react with:Bovine, Goat, Horse, Human, Sheep					
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid					
Max Ex/Em	Fluorophore	Excitation M	ax (nm)	Emission Max (nm)		
	FITC	490	. ,	525		
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant					
Buffer Solution	Phosphate buffered sa	aline				
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Alb					

Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	Porcine PBMCs		
External Database Links	UniProt: <u>Q7YRN2</u> <u>Related reagents</u> Entrez Gene: <u>397455</u> CD3E <u>Related reagents</u>		
Fusion Partners	Lymph node cells from immunised BALB/c mice were fused with cells of the NS0 myeloma cell line		
Specificity	<b>Mouse anti Pig CD3, clone PPT3</b> recognizes the porcine homologue of human CD3 $\epsilon$ , a 24 kDa single pass type I membrane protein expressed by T-lymphocytes. Clone PPT3, also known under the clone designation FY1H2, was clustered at the second international swine CD workshop and found to specifically recognise an epitope on the porcine CD3 $\epsilon$ designated as CD3c ( <u>Pescovitz, M.D., <i>et al.</i> 1998</u> ).		
	CD3 is a multimeric protein complex composed of four distinct polypeptide chains ( $\epsilon$ , $\gamma$ , $\delta$ , $\zeta$ ) that assemble and function as three pairs of dimers ( $\epsilon\gamma$ , $\epsilon\delta$ , $\zeta\zeta$ ). The CD3 complex serves as a T cell co-receptor that associates non-covalently with the T cell receptor (TCR) ( <u>Guy, C.S &amp; Vignali, D.G. 2009</u> ). CD3 is a defining feature of cells belonging to the T cell lineage, antibodies recognising pig CD3 therefore provide useful markers of porcine T cells.		
	Clone PPT3 has been demonstrated to recognise an epitope that is expressed both intracellularly and extracellularly, additionally clone PPT3 has been demonstrated to activate $\alpha/\beta$ T-cells (Kirkham P.A., <i>et al.</i> 1996).		
	Clone PPT3 was tested on PBL from a range of other mammalian species and found to be negative suggesting that the epitope recognised by this clone is specific to porcine ( <u>Yang, H. <i>et al.</i> 1996</u> ).		
References	<ol> <li>Yang, H. <i>et al.</i> (1996) Preparation of monoclonal anti-porcine CD3 antibodies and preliminary characterization of porcine T lymphocytes. <u>Immunology. 88 (4): 577-85.</u></li> <li>Kirkham, P.A. <i>et al.</i> (1996) Porcine CD3 epsilon: its characterization, expression and involvement in activation of porcine T lymphocytes. <u>Immunology. 87 (4): 616-23.</u></li> <li>Pescovitz, M.D. <i>et al.</i> (1998) Analyses of monoclonal antibodies reacting with porcine CD3: results from the Second International Swine CD Workshop. <u>Vet Immunol Immunopathol. 60: 261-8.</u></li> <li>Forberg H <i>et al.</i> (2014) Early responses of natural killer cells in pigs experimentally infected with 2009 pandemic H1N1 influenza A virus. <u>PLoS One. 9 (6): e100619.</u></li> </ol>		
Further Reading	1. Guy, C.S. & Vignali, D.A. (2009) Organization of proximal signal initiation at the TCR:CD3 complex. Immunol Rev. 32: 7-21.		

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.	
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA5951F 10041	
Regulatory	For research purposes only	

### Related Products

### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL: FITC (MCA928F)

#### **Recommended Useful Reagents**

MOUSE ANTI PIG CD4 ALPHA:RPE (MCA1749PE) MOUSE ANTI PIG wCD8 ALPHA:RPE (MCA1223PE) MOUSE ANTI PIG CD25 (MCA1736GA)

 North & South
 Tel: +1 800 265 7376
 Worldwide

 America
 Fax: +1 919 878 3751
 Email: antibody\_sales\_us@bio-rad.com

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739 Email: antibody\_sales\_uk@bio-rad.com Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368511:200529'

#### Printed on 24 Feb 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint