

Datasheet: MCA5951PE

BATCH NUMBER 155390

Description:	MOUSE ANTI PIG CD3:RPE
Specificity:	CD3 EPSILON
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	PPT3
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig		
Species Cross Reactivity	Does not react with:Bovine, Goat, Horse, Human, Sheep		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1.0 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		

Stabilisers	1% Bovine Serum Albumin 5% Sucrose
Immunogen	Porcine PBMCs
External Database Links	<p>UniProt: Q7YRN2 Related reagents</p> <p>Entrez Gene: 397455 CD3E Related reagents</p>
Fusion Partners	Lymph node cells from immunised BALB/c mice were fused with cells of the NS0 myeloma cell line
Specificity	<p>Mouse anti Pig CD3, clone PPT3 recognizes the porcine homologue of human CD3ε, a 24 kDa single pass type I membrane protein expressed by T-lymphocytes. Clone PPT3, also known under the clone designation FY1H2, was clustered at the second international swine CD workshop and found to specifically recognise an epitope on the porcine CD3ε designated as CD3c (Pescovitz, M.D., et al. 1998).</p> <p>CD3 is a multimeric protein complex composed of four distinct polypeptide chains (ε, γ, δ, ζ) that assemble and function as three pairs of dimers (εγ, εδ, ζζ). The CD3 complex serves as a T cell co-receptor that associates non-covalently with the T cell receptor (TCR) (Guy, C.S & Vignali, D.G. 2009). CD3 is a defining feature of cells belonging to the T cell lineage, antibodies recognising pig CD3 therefore provide useful markers of porcine T cells.</p> <p>Clone PPT3 has been demonstrated to recognise an epitope that is expressed both intracellularly and extracellularly, additionally clone PPT3 has been demonstrated to activate α/β T-cells (Kirkham P.A., et al. 1996).</p> <p>Clone PPT3 was tested on PBL from a range of other mammalian species and found to be negative suggesting that the epitope recognised by this clone is specific to porcine (Yang, H. et al. 1996).</p>
References	<ol style="list-style-type: none"> 1. Yang, H. <i>et al.</i> (1996) Preparation of monoclonal anti-porcine CD3 antibodies and preliminary characterization of porcine T lymphocytes. Immunology. 88 (4): 577-85. 2. Kirkham, P.A. <i>et al.</i> (1996) Porcine CD3 epsilon: its characterization, expression and involvement in activation of porcine T lymphocytes. Immunology. 87 (4): 616-23. 3. Pescovitz, M.D. <i>et al.</i> (1998) Analyses of monoclonal antibodies reacting with porcine CD3: results from the Second International Swine CD Workshop. Vet Immunol Immunopathol. 60: 261-8. 4. Forberg H <i>et al.</i> (2014) Early responses of natural killer cells in pigs experimentally infected with 2009 pandemic H1N1 influenza A virus. PLoS One. 9 (6): e100619. 5. Uehlein, S. <i>et al.</i> (2021) Human-like Response of Pig T Cells to Superagonistic Anti-CD28 Monoclonal Antibodies. J Immunol. Oct 08 [Epub ahead of print]. 6. Zhao, H. <i>et al.</i> (2022) Development of <i>RAG2</i>^{-/-} <i>IL2Rγ</i>^{-/-} immune deficient

Further Reading	1. Guy, C.S. & Vignali, D.A. (2009) Organization of proximal signal initiation at the TCR:CD3 complex. Immunol Rev. 32: 7-21. 2. Maciag, S.S. <i>et al.</i> (2022) The influence of source of porcine colostrum in development of early immune ontogeny in the piglet Ref Sq. Mar 24 [Epub ahead of print].
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA5951PE 20487
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[MOUSE ANTI PIG CD4 ALPHA:FITC \(MCA1749F\)](#)

[MOUSE ANTI PIG wCD8 ALPHA:FITC \(MCA1223F\)](#)

[MOUSE ANTI PIG CD25 \(MCA1736GA\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M375611:210104'

Printed on 22 Feb 2024